



HAL
open science

Sex Differences in Endurance Running

Thibault Besson, Robin Macchi, Jeremy Rossi, Cedric Y. M. Morio, Yoko Kunimasa, Caroline Nicol, Fabrice Vercauysen, Guillaume Y. Millet

► **To cite this version:**

Thibault Besson, Robin Macchi, Jeremy Rossi, Cedric Y. M. Morio, Yoko Kunimasa, et al.. Sex Differences in Endurance Running. *Sports Medicine*, 2022, 52 (6), pp.1235-1257. 10.1007/s40279-022-01651-w . hal-03991648

HAL Id: hal-03991648

<https://amu.hal.science/hal-03991648v1>

Submitted on 23 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sex Differences in Endurance Running

Thibault Besson¹ · Robin Macchi² · Jeremy Rossi¹ · Cédric Y. M. Morio³ · Yoko Kunimasa² · Caroline Nicol² · Fabrice Vercauysen⁴ · Guillaume Y. Millet^{1,5}

Abstract

In recent years, there has been a significant expansion in female participation in endurance (road and trail) running. The often reported sex differences in maximal oxygen uptake (VO_{2max}) are not the only differences between sexes during prolonged running. The aim of this narrative review was thus to discuss sex differences in running biomechanics, economy (both in fatigue and non-fatigue conditions), substrate utilization, muscle tissue characteristics (including ultrastructural muscle damage), neuromuscular fatigue, thermoregulation and pacing strategies. Although males and females do not differ in terms of running economy or endurance (i.e. percentage VO_{2max} sustained), sex-specificities exist in running biomechanics (e.g. females have greater non-sagittal hip and knee joint motion compared to males) that can be partly explained by anatomical (e.g. wider pelvis, larger femur-tibia angle, shorter lower limb length relative to total height in females) differences. Compared to males, females also show greater proportional area of type I fibres, are more able to use fatty acids and preserve carbo-hydrates during prolonged exercise, demonstrate a more even pacing strategy and less fatigue following endurance running exercise. These differences confer an advantage to females in ultra-endurance performance, but other factors (e.g. lower O_2 carrying capacity, greater body fat percentage) counterbalance these potential advantages, making females outperforming males a rare exception. The present literature review also highlights the lack of sex comparison in studies investigating run-ning biomechanics in fatigue conditions and during the recovery process.

Key Points

Female participation in endurance running has increased considerably over the last 20 years. Some physiological factors are known to explain sex differences in ‘classic’ endurance running performance (e.g. $\dot{V}O_{2max}$); this narrative review discusses via a multidisciplinary approach—physiological, biomechanical and neuromuscular factors that differentiate sexes in prolonged running exercise.

Although males and females demonstrate differences in running biomechanics, they do not differ in terms of running economy. A greater proportion of type I fibres and greater ability to use fatty acids in females could explain the lower neuromuscular fatigue of the quadriceps and triceps surae reported after prolonged running races compared to males, and could confer an advantage to females over ultra-endurance competitions.

More direct comparisons between males versus females in response to fatiguing running exercise are needed.

✉ Guillaume Y. Millet
guillaume.millet@univ-st-etienne.fr

¹ Univ Lyon, UJM-Saint-Etienne, Inter-University Laboratory of Human Movement Biology, EA 7424, 42023 Saint-Étienne, France

² Institute of Movement Sciences (ISM), National Center of Scientific Research (CNRS) and Aix Marseille Univ, Marseille, France

³ Decathlon SportsLab, Movement Sciences Department, Lille, France

⁴ Impact of Physical Activity on Health (IAPS), University of Toulon, Toulon, France

⁵ Institut Universitaire de France (IUF), Paris, France

1 Introduction

Running, one of the most popular recreational sports, is associated with many health benefits [1–3]. Supported by major sports companies, running as a leisure activity has grown rapidly since the 1970s in North America and in Europe since the beginning of the 1980s [3]. During the last decade, female participation in running events has increased considerably. According to a recent survey (<https://runrepeat.com/state-of-running>), covering more than 70,000 road races from 5 km to marathons all over the world, female participation had reached 50% in 2018. The International Trail Running Association recently reported an increase in female participation in trail running from 18% in 2013 to 26% in 2019 (<https://itra.run/>). In ultra-trail races, the participation rate of females is much lower, yet a sixfold increase has also been reported since the 1980s [4]. Despite the increasing female participation, the vast majority of scientific studies are still conducted on males. Indeed, based on data collected from three major sport and exercise medicine journals, the average ratio of male to female participants per article is ~ 65:35 [5]. Part of this difference has been attributed to sex differences in willingness to participate in certain types of research. In other words, a volunteer bias in female participation in running research may exist [6].

With the growth of the number of females engaged in running events, their performance rapidly improved through the 1980s [7] and, from then, researchers started to compare male and female running performance from sprint to marathon distances [8, 9]. In the 1990s, it was believed that there would be a reduction [10–12] or even disappearance [8] of sex differences in running performance. For instance, using a regression analysis between running velocities over several distances and historical time in males and females, Whipp and Ward [8] reported in 1992 that females could outrun males in marathon distance in 1998. However, the difference in performance between sexes remains around 10–12% for running events from sprint to marathon [13–15]. The issue of sex differences in performance as a function of distance is complex. When using the world best times from 100 m to 200 km, Coast et al. [14] reported that the difference in performance increases with distance, particularly in events longer than the marathon. It has also been reported that the difference between sexes when comparing the best male and the best female runners was ~ 17% for the 100-km Lauf Biel in Switzerland [16] and ~ 20% for the 161 km of various north American ultra-marathons [7]. Additionally, da Fonseca-Engelhardt et al. [17] observed the same tendency on two of the hardest ultramarathons worldwide, i.e. the ‘Badwater’ in the USA and the ‘Spartathlon’ in Greece, with performances being ~ 20% slower for females than for males (based on the top five finishers). On the contrary, Peter et al.

[18] reported a ~ 10–13% gap when comparing the fastest male and female runners ever of the 24-h ultra-marathon. How do we explain the variability in sex differences in running performance between studies regarding events longer than the marathon? First, reported sex differences depend on how the question is framed, for example, comparing the top female and male runners or considering all participants. In the latter case, sex differences seem to even decrease with increasing race distance and no differences were reported in races longer than 315 km (e.g. <https://runrepeat.com/state-of-ultra-running>), whereas, based on unpublished data, the sex difference in performance appears constant with increasing distance when considering the first males and the first females. Second, the percentage of female participation in these events is much lower than reported for males, being as low as 10% in some events, which may confound the performance differences in the very long-distance races [19]. Third, many of the races above the marathon distance are trail running events. These races are not as standardized as road races, and consist of regular changes in terrain, technical difficulty, elevation gain and exact distance, and it is not known if this has a greater impact on males than females, or vice versa. Despite the substantially lower participation rate of females, ultra-distance in running is one of the rare disciplines where females are able to outperform men. To cite only a few, Corinne Favre, Pamela Reed, Hiroko Okiyama and Jasmin Paris finished at first place overall at the ‘CCC’ in 2006 (86 km), ‘Badwater’ 2002 and 2003 (217 km), the ‘Deutschland-lauf’ in 2007 (1,204 km), and the 268-mile Montane Spine race, respectively. Jasmin Paris not only won the race, but also shattered the previous male record by 12 h. It is, however, likely that these exceptional performances are not only due to the great mental and physical capacities of these runners but also to the fact that the best men were not running in these particular events. In summary, although the question of the evolution of performance across race distance according to sex is a complex one, largely due to a question of parity in trail and ultra-trail running participation, the sex gap in performance appears relatively constant with increasing distance when focusing on top level runners.

Long-distance running performance (up to the marathon) depends on the interaction between $\dot{V}O_{2\max}$, the fraction of $\dot{V}O_{2\max}$ sustained (F) on the covered distance that is closely linked to lactate threshold (LT), and running economy (RE) (e.g. [20, 21]). Endurance factors such as F or LT do not seem to be affected by sex (e.g. [22–25]). For instance, Davies and Thompson [22] reported that male and female athletes were able to sustain similar F over the marathon distance (82 and 79%, respectively). The most obvious physiological variable to explain running performance difference is the maximal oxygen uptake ($\dot{V}O_{2\max}$). When expressed in $\text{mLO}_2 \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$ (of body mass), $\dot{V}O_{2\max}$ is considered as the main factors explaining sex differences in endurance

running performance [20, 26], mainly due to greater percentage of body fat and lower haematocrit levels in females [26, 27]. When $\dot{V}O_{2\max}$ in males and females is adjusted to fat-free mass, some controversy exists in the literature, with some studies showing no sex differences [28–30] whereas other report a higher $\dot{V}O_{2\max}$ in males [31, 32]. Lower O_2 -carrying capacity is definitively one factor explaining the lower $\dot{V}O_{2\max}$ in females [33]. As sex differences in $\dot{V}O_{2\max}$ have largely been reviewed elsewhere [26, 31, 32, 34], the present review focuses on the other factors that differentiate males and females in endurance running. Thus, the aim of the present narrative review was to examine whether males and females differ in terms of running biomechanics, economy and physiological responses during short (< 1 h) and prolonged (i.e. several hours of running) running exercises. This narrative review offers a novel comprehensive and multidisciplinary (i.e. psychological, physiological, neuromuscular and biomechanical aspects) overview of the sex differences in endurance running. Article databases of the US National Library of Medicine (PubMed), ScienceDirect and SPORTDiscuss were searched using the search general (Gender or Sex differences) and specific terms (corresponding to the studied scientific area of each part). For instance, article databases focusing on the “Running economy” part were searched as follows: Gender OR Sex differences AND Running economy OR oxygen demand OR Energy cost of running OR Oxygen cost of running OR oxygen uptake OR metabolic efficiency. Any additional relevant literature was obtained from the reference lists of the published papers. This searching process has been applied for each part of the current review.

2 Biomechanics of Running

Many studies have been conducted to compare the running biomechanics of males and females (Table 1). Several parameters such as the studied population, footwear, running surface and speed may contribute to the disparity in the results observed between studies. In the absence of fatigue, differences in spatiotemporal, kinematic and kinetics parameters of the running pattern have been extensively studied, but primarily in experimental conditions at given absolute running speeds and for a short period of time rather than in endurance condition. The major sex differences are indicated in blue on Fig. 1.

2.1 Spatiotemporal Parameters

Sex differences in spatiotemporal parameters have been investigated over a large range of running velocities (from 7.5 to 24 km h⁻¹) [35–38]. For given absolute running

speeds above 16 km h⁻¹, female runners present shorter absolute contact time, longer absolute flight time, shorter absolute stride length and higher stride frequency than males [36]. Because of shorter contact time and longer flight time, females present a lower duty factor, i.e. percentage of stride time spent on the ground, indicating that females are not a “scaled-down version of the male model” [36]. Interestingly, these sex differences seem to disappear at slower speeds (10–14 km h⁻¹), with the exception of contact time as females still present shorter contact time than males when the speed decreases [35]. At high absolute speeds, females probably have to adjust their biomechanics, in particular through adopting a high stride frequency as they are closer than males to their maximal speed. Indeed, once normalized to maximal velocity (from 60 to 90% of maximal velocity), sex differences in contact time and in duty factor tend to disappear and step frequency is actually lower in females [36]. Sex differences in body proportions might also explain part of the sex differences in spatiotemporal parameters. For instance, several studies reported shorter step length in females at various running speeds (from 7.5 to 24 km h⁻¹), but these differences vanished or even reversed once adjusted to stature or leg length [35, 36, 38, 39]. Furthermore, a significant positive association between contact time and body height, leg length and body mass has been found [37], but the extent to which morphological differences contribute to the sex influence on spatiotemporal parameters is not yet clear.

2.2 Running Kinematics and Muscle Electromyographic Activity

The multitude of variables studied and the diversity of the methods of running kinematics analysis make comparisons between studies quite challenging. However, a shared observation across studies is that sex differences mostly existed at the hip and knee level [40–51].

In recreational runners, females exhibited significantly greater hip adduction and internal rotation during the stance phase of running compared to males across a range of speeds [40, 42, 43, 49] and surface inclinations [42]. Sex differences in running kinematics at the hip level have been attributed to the fact that females have a wider pelvis [52] and possibly weaker hip abductor and external rotator muscle strength compared to males [49, 53]. Despite moderate correlations between hip, pelvis and trunk kinematics and hip abductors or external rotator strength in both males or females, hip strength parameters remain a substantial portion of the explained variance in running kinematics for both sexes [45]. Females have also been shown to present greater knee abduction during the stance phase of running compared to males [43, 44, 47–50]. Greater knee abduction has been associated with smaller rearfoot

Table 1 Overview of sex differences in running biomechanics

References	<i>n</i>	Footstrike	Footwear	Surface	Speed (km h ⁻¹) and slope	Spatiotemporal parameters	Kinematics	Kinetics
Almonroeder and Benson [40]	14 M 18 F	Rearfoot	Std	↘	14.4		<p><u>Peak angles:</u> Knee flex : = Knee add: = Knee IR: = Hip add: F+ Hip IR: F+</p>	<p>MOMENT Peak moment: Knee ext moment: F- PFJ stress: F- PFJ reaction Force: F-</p>
Bazuelo-Ruiz et al. [41]	28 M 27 F	Rearfoot	Std	↘	11.9		<p><u>Peak angles:</u> Knee flex: F+ Ankle dorsiflexion: F+</p>	<p>GRF Impact peak: = Active peak: F- Loading rate: F+ Peak braking force: = Peak propulsive force: F+</p>
Chumanov et al. [42]	17 M 17 F	NS	NS	↘	6.5, 9.7, 13 With 0, 10 and 15% uphill incline		<p>For all speed and incline: <u>Peak angles:</u> Hip add: F+ Hip IR: F+ Lateral pelvic tilt: = <u>Amplitudes:</u> Hip add: F+ Hip rotation and lateral pelvic tilt: =</p>	
Esculier et al. [193]	51 M 36 F	NS	Std	↘	Preferred running speed* M: 11.2 F: 10.1	<p>Stride length: F- Step width: F-</p>	<p><u>Peak angles:</u> Knee flex: = Knee add: = Knee abd: =</p>	<p>MOMENT Peak moment: Medial TFJ forces: F- External knee flex: F- External knee add: =</p>
Ferber et al. [43]	20 M 20 F	Rearfoot	NS	↘	13.1	Contact time: =	<p><u>Peak angles:</u> Hip flex: = Hip add: F+ Hip IR: F+ Knee flex: = Knee abd: F+ Knee IR: =</p>	<p>MOMENT Peak moment: Hip ext, add and IR: = Knee ext, abd and IR: = WORK Negative work: Hip sagittal plane: = Hip frontal and trans planes: F+ Knee sagittal, frontal and trans planes: =</p>

Table 1 (continued)

References	<i>n</i>	Footstrike	Footwear	Surface	Speed (km h ⁻¹) and slope	Spatiotemporal parameters	Kinematics	Kinetics
García-Phillos et al. [35]	51 M 46 F	NS	NS	⚡	10, 11, 12, 13, 14, 15, 16	Contact time: F- at 10 to 16 km h ⁻¹ Flight time: = at 10 to 16 km h ⁻¹ Stride length: = at 10 to 13 km h ⁻¹ F- at 14 to 16 km h ⁻¹ Stride length (norm.): = at 10 to 14 km h ⁻¹ F- at 15 and 16 km.h ⁻¹ Step frequency: = at 10 to 13 km h ⁻¹ F+ at 14 to 16 km h ⁻¹	Peak angles: Hip add: F+ Knee add: F-	MOMENT Peak moment: Hip add at Impact and Mid-stance: F+ Knee add at Impact and Mid-stance: =
Gehring et al. [44]	16 M 16 F	Rearfoot	Std	⚡	10.8, 14.4, 18	Contact time: F-	Peak angles: Hip add: F+ Knee add: F-	WORK Hip joint impulse: F+ Knee joint impulse: F+
Hannigan et al. [45]	37 M 23 F	NS	Std	⚡	Self-selected "easy" run pace* M: 12.6 F: 11.9		Amplitudes: Hip ext: F+ Hip add: = Hip IR: F+ Anterior pelvic tilt, Contralateral pelvic drop and Pelvic IR: =	
Hennig [194]	17 M 15 F	NS	Std	⚡	11.9	Contact time: =		GRF Impact peak: F- Active peak: =

Table 1 (continued)

References	<i>n</i>	Footstrike	Footwear	Surface	Speed (km h ⁻¹) and slope	Spatiotemporal parameters	Kinematics	Kinetics
Isherwood et al. [60]	20 M 20 F	Rearfoot	Std	▲	11.9	Contact time: F-	Peak angles: Hip flex and add: = Hip IR: F+ Knee flex, add: = Knee IR: F- Ankle eversion: = Ankle add: =	GRF Impact peak and active peak: = Loading rate: F+ Peak braking force and propulsive force: = MOMENT Peak moment: In the sagittal plane, for Hip, Knee and Ankle: F-
Keller et al. [63]	13 M 10 F	NS	Std	▲	From 12.6 to maximal speed			GRF Active peak: F- at 18 km h ⁻¹ Loading rate: =
Malinzak et al. [46]	11 M 9 F	NS	Perso	▲	M: 18.8 F: 18.4		Peak angles: Knee flex : F-	
Nelson et al. [36]	24 M 21 F	NS	NS	▲	16.5 to 24.1	Absolute speed: Stride length, contact time and duty factor: F- Step frequency and flight time: F+ Stride length (normalized): F+ Relative speed: Stride length, step frequency and duty factor: F- Contact time: = Flight time: F+ Stride length (normalized): F-		
Nigg et al. [47]	47 M 46 F	Rearfoot	Std	▲	12		Amplitudes: Pelvic tilt: F+ Hip add: F+ Knee abd: F+	
Phinyomark et al. [48]	220 M 263 F	NS	Std	▲	Self-selected speed 8 to 12.1		Peak angles: Hip add: F+ Knee abd and ER: F+ Knee flex: F- Ankle eversion and dorsiflexion: F-	
Roche-Seruerdo et al. [37]	52 M 45 F	NS	Perso	▲	12	Contact time: = Flight time: = Step length: = Step frequency: =		

Table 1 (continued)

References	<i>n</i>	Footstrike	Footwear	Surface	Speed (km h ⁻¹) and slope	Spatiotemporal parameters	Kinematics	Kinetics
Sakaguchi et al. [49]	11 M 11 F	NS	Std	$\underline{\text{A}}$	12.6	Contact time: =	Peak angles: Hip add and IR: F+ Knee IR: = Knee abd: F+ Tibial IR: = Rearfoot eversion: F-	
Schache et al. [39]	22 M 22 F	NS	Perso	$\underline{\text{A}}$	14.4	Stride time: F- Stride frequency: F+ Stride length: F- Stride length (normalized): = Contact time: F- Flight time: F- Contact time: =	Amplitudes: Pelvis (antero post tilt, obliquity and axial rotation): F+ Hip flex-ext and IR-ER: = Hip Add-abd: F+	
Sinclair et al. [50]	12 M 12 F	NS	Std	$\underline{\text{A}}$	14.4	Contact time: =	Only significant differences are reported Peak angles: Hip flex: F- Knee abd and IR: F+ Ankle eversion: F+	GRF Vertical impact peak : = Loading rate: = Peak braking force and prop force: = Peak medial force and lateral force: =
Sinclair and Selfe [67]	15 M 15 F	Rearfoot	Std	$\underline{\text{A}}$	14.4			MOMENT Peak moment: Knee extensor and abd: F+ PTJ contact force: F+ PTJ load rate: F+
Sinclair and Taylor [55]	20 M 20 F	NS	Std	$\underline{\text{A}}$	14.4		Peak angles: Ankle eversion: F+ Tibial IR: F+ Amplitudes: Ankle eversion and Tibial IR: =	

Table 1 (continued)

References	<i>n</i>	Footstrike	Footwear	Surface	Speed (km h ⁻¹) and slope	Spatiotemporal parameters	Kinematics	Kinetics
Takabayashi et al. [38]	12 M 12 F	Rearfoot	Barefoot	$\dot{\alpha}_s$	Individual transition speed from walking to running on a treadmill* M: 7.8 F: 7.3	Relative speed: Step frequency: = Step length: F- Step length (normalized): =	Only significant differences are reported Peak angles: Rearfoot plantarflexion: F+ Midfoot dorsiflexion and Abd: F+ Amplitudes: Rearfoot sagittal plane: F+ Midfoot sagittal plane: F+	
Willson et al. [51]	19 M 19 F	NS	Std	$\dot{\alpha}_s$	Between 12.7 and 14		Amplitudes: Hip add and ER: = Knee IR: F- Knee add: = Angle at mid stance: Hip add: F+ Hip ER: = Knee IR and add: =	

For kinematic variables, only peak and amplitude joint angles were reported. The results reported are based on the *p* value according to the significant level set in each studies

M males, *F* females, $\dot{\alpha}_s$ overground, $\dot{\alpha}_s$ treadmill, *GRF* ground reaction forces, *Std* standardized footwear (i.e. all participants wore the same pair of shoes), *Perso* personal shoes, *Flex* flexion, *Ext* extension, *Abd* abduction, *Add* adduction, *IR* internal rotation, *ER* external rotation, *PFJ* tibio-femoral joint, *PFJ* patello-femoral joint, *NS* not specified, * significant sex difference, *F+* superior in females, *F-* inferior in females, = no sex difference




Fig. 1 This schematic representation of a female runner is an overview of the main psychological, physiological, neuromuscular and biomechanical sex differences in endurance running. Parameters that could give an advantage to males and to females in endurance run-

ning performance are boxed in green and orange, respectively. >: superior in females, <: inferior in females, =: no sex difference, AT Achilles tendon, O_2 oxygen, $\dot{V}O_{2max}$ maximal oxygen uptake

eversion to compensate for larger hip adduction, as typically seen in females [49], i.e. sex differences in kinematics at the hip level can explain differences at the knee level. In other words, greater knee abduction in females could also be a plausible consequence of morphological differences and especially the greater Q-angle (i.e. the acute angle formed by the vector for the combined pull of the quadriceps femoris muscle and the patellar tendon [54]) reported in females [43, 50]. Smaller knee flexion angles have been reported in females both at high running velocities (around 18 km h^{-1} [46]) and at preferred running speed among a very large sample of 483 (263 females and 220 males) recreational and competitive runners [48]. However, previous investigations showed no sex difference in knee sagittal plane motion between females and males [43, 50]. These discrepancies could be explained by the imposed speed (13 and 14.4 km h^{-1}) in previous studies [43, 50], where running speeds varied from 8 to 12 km h^{-1} in Phinyomark et al. [48]. At the ankle level, one study found smaller rearfoot eversion in female runners [49], whereas two other studies found greater rearfoot eversion in females [50, 55]. In these studies, as males and females ran at the same absolute speed, Takabayashi et al. [38] suggested that differences could be attributed

to differences in body size and spatiotemporal parameters [38], as described in Sect. 2.1.

Given the identified sex differences in joint kinematics, research has been conducted to identify sex specificities in lower limb muscle activities. Unfortunately, sex comparisons have mostly been studied at given absolute running speeds and for very short periods of time (about 10 s). The neuromuscular strategies adopted may thus differ from those used in endurance running and racing conditions. In these experimental running conditions, females had a higher activation of the gluteus maximus [42, 51] than males at all speed (from 9.7 to 14 km h^{-1}) and incline (0 , 10 and 15%) conditions and a larger increase in activity of the vastus lateralis than males as incline increased [42]. Females may thus use different neuromuscular strategies than males in case of increased task demand, i.e. changes in speed and incline [56]. At similar running speeds of around 18 km h^{-1} , Malinzak et al. [46] demonstrated a higher activation of the quadriceps but a lower activation of the hamstrings in recreational female athletes compared to males, which potentially induced a higher anterior knee shear force (i.e. perpendicular force applied by the quadriceps muscles on the tibia) in female runners. In contrast, a higher medial hamstring pre-activation has been reported

in downhill running at a slow running speed (7.5 km h^{-1} on a 15° declined treadmill) in female runners, which may contribute to better stabilization of the knee joint in this situation [57]. Similarly, an increased pre-activation of the peroneal muscle group was found in female runners during level running, and this was considered as a strategy to stabilize and protect the ankle joint [58]. As these studies only considered a small number of muscles, further analysis of the inter-muscular coordination in running was limited. However, in a recent study conducted in 60 females and 60 males [59], the concept of muscle synergies could be used based on the electromyographic recording of 13 lower limb muscles at running speeds ranging from 7.2 to 13.7 km h^{-1} . This study found sparse sex-specific modulations of the relative muscle contribution in running. During the stance phase, no sex differences were found in the contribution of the hip extensors, tensor fasciae latae and plantar flexors, yet females used the vastus medialis less during the weight acceptance. In this particular phase, the timing of the main peak of activity was similar but the synergy was narrower in females, implying a shorter time-duration of activation in females than in males. The authors suggested that the neuromotor strategy showed by females could reduce their fatigability during prolonged locomotion efforts. During the early swing, females used the rectus femoris less but the gluteus medius more whereas during the late swing, females used the tibialis anterior more compared to males.

Overall, despite sex-specificities of the running kinematics, the reported differences in muscle activities, such as the female-specific protective joint strategies, are still controversial and unlikely to reduce the sex-related gap in performance. Further studies should focus on the sex differences in muscle synergies under endurance and competitive running conditions.

Most of these studies investigated running mechanics at the same absolute speed, putting female runners at a greater relative demand on their musculoskeletal system. Indeed, most of the previous literature on kinematics differences between male and female runners failed to establish a relationship with sex, often because of the lack of control for confounding factors such as running speed and participants' height and body mass. It is therefore necessary to conduct more studies with larger sample sizes allowing consideration of confounding factors in order to obtain a clearer picture of the real influence of sex on running biomechanics.

A recent study assessed sex differences in running biomechanics in a population of Chinese males and females [60]. The authors report similarities with sex differences in Western populations (e.g. increased hip and knee range of motion in females), but also differences in biomechanics (e.g. knee and ankle joint kinematics, vertical ground

reaction force and plantar pressure) between Western and Chinese runners. Thus, the authors emphasized the fact that ethnicity should be considered when comparing running biomechanics in males and females, in particular by running shoe manufacturers. Another under-studied topic is the influence of females' breasts on their running biomechanics. A recent review [61] pointed out that, although there is some information on breast motion during walking or running, the previous breast-focused literature had "poor research design and inadequate biomechanical methods used to quantify complex three-dimensional breast motion". Brown and Scurr [62] found a slightly negative influence of breast mass on the marathon finish-time; however, no studies have been conducted so far to determine the influence of the biomechanics of females' breasts and trunk on their running performance. This specificity of females' morphology should also be considered in future studies examining sex difference in running biomechanics.

2.3 Running Kinetics and Muscle–Tendon Stiffness Regulation

At given running speeds up to 21.6 km h^{-1} , Keller et al. [63] did not report any sex difference in the changes in maximal vertical ground reaction force and loading rate (normalized by body weight) with speed. However, sex differences in the normalized force data have been reported in a larger group of recreational runners when running at the same absolute running speed of 11.9 km h^{-1} [41]. Females exhibited a significantly higher loading rate but a lower active peak force than males in both pre-fatigue and fatigue conditions. They also showed higher peak propulsive force (antero-posterior), but only before fatigue. Again, as this study was conducted at the same absolute running speed, greater peak propulsion force (measured on the antero-posterior axis) observed in females may have been due to the greater relative effort compared to males.

Joint moments and mechanical stress applied to lower limb joints have been compared to better understand potential sex differences in running-related injuries, rather than running performance [64]. Specifically, females appeared to be at greater prevalence [65] and incidence [66] to sustain patellofemoral joint pain compared to males. Conflicting results exist on sex differences in patellofemoral joint mechanical stress (PFJS) [40, 67, 68], probably due to different methods used to calculate it. In addition, contradictory results exist in sex differences when considering peak knee extension moment (normalized by body mass) in running since no difference [43], greater [67] or lower [40] moment in females have been reported. In the frontal plane, females have been shown to display greater normalized hip and knee adduction moment compared to males [44]. Similar to sex

differences in running kinematics, sex differences in kinetics have mainly been attributed to – yet not measured—morphological sex differences.

Running is a stretch–shortening cycle type activity in which higher Achilles tendon (AT) stiffness is expected to affect the storage of elastic energy during the braking phase, followed by its recoil during the subsequent push-off phase. Therefore AT stiffness is expected to affect RE [69–73]. Tendon structural quality and resistance to deformation, tendon collagen synthesis rate and hypertrophic effect of regular running exercise on the patellar and AT have been reported to be lower in females than in males [74, 75]. In particular, no significant hypertrophic effect on the patellar and AT was found in young and regularly trained female runners (at least 40 km of running mileage per week for the previous 5 years) as compared to untrained ones, whereas a significant gain was found in similarly trained males [75]. Both non-athletic [71] and trained female runners [73] have been reported with lower stiffness and hysteresis of AT than their male counterparts. Despite this AT difference, trained female and male runners had a similar RE (see Sect. 2.2), but this remains the only study on such a sex comparison. Other confounding factors should be considered, such as AT length and moment arm, but these factors were studied primarily in elite male runners [76]. Thus, further work is needed to determine in females the exact influence of a less stiff AT on their RE and running performance.

In addition to their effects on tendons, oestrogens are reported to reduce ligament stiffness. Although suggested as a potential risk factor for anterior cruciate ligament (ACL) injuries for female participants during sports activity [77], a causal link is refuted by other studies reporting no associated decrease in anterior knee stiffness [78]. ACL laxity also appears to be affected by the menstrual cycle, as evidenced by its increase reported in the late follicular phase, i.e. when the serum level of estradiol peaks and progesterone is low (e.g. [78, 79]). Once again, no associated change was found in anterior knee stiffness [78]. Within this framework, muscle stiffness being an important component of joint stability, it is noteworthy that Khowailed et al. [80] revealed different muscle activation strategies in running during different phases of the menstrual cycle. During the follicular phase, when oestrogen levels are low, females placed greater reliance on the quadriceps than on the hamstrings. During ovulation, when oestrogen level and knee laxity are high, increased hamstring pre-activation and co-activation during the braking phase have been observed. Thus, as oestrogen levels seem to influence the stretch–shortening cycle and potentially RE, running performance could possibly vary across the menstrual cycle.

3 Substrate Utilization and Running Economy

3.1 Substrate Utilization

Numerous original studies and review studies have analyzed sex differences in substrate utilization during exercise lasting from 60 to 120 min (e.g. [81–86]) at different intensities (from 35 to 75% $\dot{V}O_{2max}$). Comparing sexes in substrate use is methodologically challenging since this parameter is influenced by several factors such as training status, menstrual cycle phase, diet, age, etc. Controlling for age, diet and physical activity level (i.e. using $\dot{V}O_{2max}$ expressed relative to fat-free mass) would allow avoiding heterogeneity in the results when comparing exercise-induced metabolic responses between males and females [82]. Menstrual cycle phase has been reported to have a small impact on sex difference in substrate utilization [86]. Oral contraceptives have a slightly greater impact on substrate oxidation [86]. Together this implies that when the variance of these two factors is not controlled, sex differences in substrate utilization may be biased.

When considering only the studies that presented as well controlled with regard to these factors, the overall conclusion is that females oxidize more lipid and less carbohydrate (CHO) and protein than males during prolonged exercise (at least until 120 min of exercise duration). Females have also greater metabolic flexibility since substrate oxidation is readily adjusted according to the availability of nutrients. In this regard, the respiratory exchange ratio (RER) has been reported to be lower in females during low-intensity, prolonged exercises, supporting a lower reliance on whole-body CHO oxidation (e.g. [86]). For instance, lower RER values were found for females (0.87) than for males (0.94) during a 15.5 km treadmill run at 65% VO_{2max} [84]. During an incremental treadmill run test to exhaustion performed in a large cohort of 157 males and 143 females, Venables et al. [87] showed that females displayed higher maximal rates of fat oxidation. In addition, females exercised at a higher % VO_{2max} , for which a CHO as dominant fuel is observed. Although less used than CHO and fat, protein provides approximately 5–8% of total energy in both sexes [29] during prolonged exercise. However, it should be noted that the protein catabolism seems to be lower in females, potentially due to the higher fatty acid oxidation [29, 30, 84, 88, 89]. Theoretically, this may constitute an advantage for females during ultra-endurance exercises (Fig. 1).

The higher metabolic flexibility observed in females constitutes a sex-specific difference that depends on several factors (and their interaction), including higher skeletal muscle insulin sensitivity as well as high levels of circulating and intramuscular fatty acids required for the oxidation

process (for a review, see [83]). Some authors have reported the crucial role of female sex steroid hormones (e.g. oestrogens) in the regulation of whole-body insulin sensitivity and substrate metabolism (e.g. [83, 86, 90]) at rest and during endurance exercise. Within the endogenous forms of oestrogens, 17- β -estradiol and associated oestrogen receptors α are important in the regulation of substrate metabolism in skeletal muscle. Higher levels of 17- β -estradiol are reported to spare glycogen stores and decrease liver glucose output and muscle uptake by shifting metabolism toward free fatty acids at certain (low to moderate) exercise intensities [28, 86, 91–93]. This metabolic hormonal action might thus contribute to enhanced performance capabilities of females compared to males for ultra-endurance exercise. Although lower RER values have been reported at submaximal running intensities (30–60% VO_{2max}) in the luteal phase of the menstrual cycle [94, 95], other studies reported no significant differences throughout the menstrual cycle (e.g. [96]). Part of this discrepancy has been attributed to the complex balance between oestrogen and progesterone concentrations and their fluctuations during the menstrual cycle [82]. According to D'Eon et al. [92], the ratio of oestrogen/progesterone would be sufficiently elevated to lead to metabolic changes between the menstrual phases.

Although females and males do not differ in their glycogen stores and capacity for storage of glucose into glycogen [83, 84], females have a higher plasma fatty acid availability due to a greater number of intramyocellular lipid (IMCL) droplets (and not larger droplets) [97]. Moreover, female skeletal muscle is reported to be better able to store lipids and oxidize fatty acids during exercise, which would contribute to maintaining a high level of IMCL turnover [83]. During exercise, this dependence on lipid substrate to cover energy expenditure in females might be explained by a greater relative area of type I muscle fibers (see next section) characterized by higher oxidative capacity, capillary density and higher insulin sensitivity [98]. Indeed, Essén et al. [99] previously reported that IMCL content was 2.8-fold higher in type I fibers compared to type II fibers. Despite inconsistencies in the literature [82], higher IMCL use during submaximal exercise has been reported in females as compared to males, irrespective of training status [100–102].

The majority of studies focusing on sex differences in substrate oxidation have been conducted on a cycle ergometer rather than in running (see Table 1 in Tarnopolsky [86]). The differences in substrate oxidation observed between sexes highlights the importance of quantifying the cycle menstrual phase, oral contraception, training level and diet in females.

3.2 Running Economy

RE is known to be the most predictive variable of running performance within homogeneous groups of trained athletes (i.e. similar VO_{2max} values) (e.g. [21]). RE depends on several physiological, biomechanical, anatomical and neuromuscular (NM) attributes, supporting the importance of a holistic approach in RE determination (for reviews, see [103, 104]). Oxygen cost in $mlO_2 \text{ kg}^{-1} \text{ km}^{-1}$ is derived from the mass-specific VO_2 , dividing the steady-state VO_2 the resting value by the running speed. Given that substrate metabolized to provide energy may vary with exercise constraints such as intensity [105], duration [106], training and sex (e.g. [97, 107]), it has been proposed to express RE as gross energy cost [105] in $J \cdot \text{kg}^{-1} \cdot \text{m}^{-1}$ or $\text{kcal} \cdot \text{kg}^{-1} \cdot \text{km}^{-1}$. An enhanced/improved RE is associated with a decrease in this value.

During submaximal running tests (< 10 min), no scientific consensus exists on the differences in RE between trained or recreational male and female runners, with conflicting results strongly related to differences in methodological setting [73, 108–114]. VO_{2max} values, training background, running performance, body fat and quantification of RE have probably contributed to the lack of consensus on this issue. In this regard, sex-specific differences have often been observed when authors expressed RE from oxygen cost or submaximal VO_2 (e.g. [23, 108, 112, 115]). Moreover, the majority of studies investigating sex differences in RE have used small sample sizes (e.g. $n < 30$) and/or comparisons across absolute speeds (rarely with relative intensities). In a study comparing RE between trained female and male runners, Fletcher et al. [73] did not show any sex difference at similar relative intensities (75, 85 and 95% of the speed at LT) when expressed in either energy cost or oxygen cost, or when using of an allometric scaling factor in order to take into account sex differences in body mass. These findings have been confirmed by the same research group with the inclusion of elite and trained male and female runners [113]. In addition, using a larger sample size ($n=95$), Black et al. [109] recently reported no sex differences in RE in $\text{kcal} \cdot \text{kg}^{-1} \cdot \text{km}^{-1}$ at similar relative and absolute running speeds among females and males of equivalent standard. Considering these findings, no clear difference in RE responses exists between male and female runners even when applying different RE units.

In most of the articles focusing on RE responses in females, a qualitative analysis of menstrual cycle is lacking, and this might constitute an important methodological bias in RE evaluation. During endurance exercises, females generally use less CHO and have lower RER values [107, 116, 117]. These metabolic changes may depend on the phases of the menstrual cycle as well as short-term oral contraceptives and, in turn, affect the RE response in females. RE may also be naturally influenced by the specificity of each phase (e.g.

early, late follicular and mid-luteal phases), and, therefore, circulating hormone levels (i.e. oestrogen and progesterone). For instance, investigating three different times during the menstrual cycle in females taking oral contraceptives (i.e. constant doses of oestrogen and progesterone for 21 days), Giacomoni and Falgairette [118] reported an improved RE at low intensities (7–9 km h⁻¹) only during the late period of oral contraceptive use (i.e. days 19–21). In this regard, Dokumacı and Hazır [119] recently reported an improved RE (i.e. in terms of energy or oxygen cost) in the luteal phase compared to the follicular phase in female competitive athletes with no oral contraceptives. Conversely, two studies have indicated that the mid-luteal phase (high levels of oestrogen and progesterone) was associated with worse RE responses compared to early and late follicular phases [120, 121]. More work is needed to better understand the effects of the menstrual phase in RE, but these findings suggest that those effects must be considered.

4 Fatigue Resistance

NM function, RE and biomechanics are often measured in a non-fatigued state. However, our group and others have consistently shown in the last two decades using laboratory or field experiments (e.g. marathon, trail/ultra-trails), that fatiguing endurance running exercise drastically affects maximal strength, RE and running patterns. Some sex differences exist at these levels and are discussed below.

4.1 Neuromuscular Fatigue

NM fatigue has traditionally been defined as an exercise-induced reduction of maximal strength or power [122]. The causes can be peripheral (i.e., within the muscle) and/or central (i.e., proximal to the neuromuscular junction), and NM fatigue etiology in running depends on many factors such as exercise duration, intensity, altitude, and positive and negative slope [123]. Central fatigue reflects an inability to voluntarily activate muscles, and has been described as a protective mechanism acting to limit peripheral perturbations [124, 125]. In contrast to central fatigue, which occurs predominantly after very long and low intensity exercise, peripheral fatigue is greater at higher intensities, shorter duration exercise [126] including downhill running [127], and could impair (1) action potential transmission along the sarcolemma, (2) excitation–contraction coupling, and (3) actin–myosin interaction [123].

Sex differences in fatigability have been mostly investigated in laboratory conditions (for a review, see [128]). While females are less strong and less powerful than males, they were found to be less fatigable than males for sustained or intermittent isometric contractions performed at a similar

relative intensity [129]. However, sex differences in fatigability are less clear following dynamic tasks as females have been reported to be either less than or as fatigable as males (see Fig. 1 in Hunter [128]). Among these studies, a lower fatigue index (i.e. smaller decline in each isokinetic variable) was found in females compared to males after a series of 30 maximal dynamic knee extension [130]. This finding could be of importance given that in a recent paper, Ehrstrom et al. [131] showed local endurance of the knee extensors was a good predictor of performance on a short trail running exercise (27 km with 1400 m of positive elevation) in trained male runners. On a fatiguing cycling task, Ansdell et al. [132] compared males and females after normalizing exercise intensity to critical intensity, and found that females displayed a greater relative critical intensity compared to males, yet contrary to their hypothesis, females had a longer time to task failure than males above critical power [132]. Females also exhibited lower strength loss and peripheral fatigue at exhaustion. However, sex differences in fatigability following prolonged running exercise is not well documented. After 2 h of laboratory treadmill running at a speed corresponding to their first ventilatory threshold, Glace et al. [133] reported that knee extensor/flexor strength measured in isokinetic conditions (60° s⁻¹) decreased in males only, while decreased hip strength was independent of sex.

Under real endurance racing conditions on short distances (<40 km), Boccia et al. [134] reported no sex difference in the decrease in knee extension force after a half-marathon performed by moderately trained runners under real racing conditions. These authors also found that the origin of fatigue was independent of sex. After a graded 20-km race (\pm 350 m) in recreational males and female runners, sex differences were found in both the acute and delayed (up to 4 days) recovery period [135]. Females showed smaller functional decrements within the first 2 h and earlier full recovery than males. Indeed, only males showed decreases in maximal isometric knee extension force and maximal power in drop jump up to 4 days after the race. However, regarding the relative decrements (normalized by pre-values), no sex difference was found, so that the lower fatigue effects observed in females may be partly attributed to their lower initial strength levels.

Over a much longer distance, Temesi et al. [136] assessed NM fatigue in experienced male and female ultra-endurance trail runners matched by relative level of performance (i.e. same average percentage of performance as the winner of each sex) after a 110-km ultra-trail-running. In the ultra-trail study, female runners had (1) less peripheral fatigue in the plantar flexors and (2) a lower decrease in maximal force in the knee extensors when compared to their male counterparts. The authors suggested that females were more fatigue resistant following ultra-endurance running effort, which could partly explain why they manage to outperform males

in races of this format. A recent study from our group investigated sex differences in NM fatigue in trail running races of various distances (from 40 to 170 km) [137]. Male and female participants were matched by a relative level of performance using the same criteria as in Temesi et al. [136]. Greater knee extensor strength loss was found in males compared to females and, contrary to our hypothesis, this sex difference was independent of the race distance. Interestingly, we observed greater peripheral fatigue on plantar flexor muscles in males compared to females in races under 60 km but no sex difference in races above 100 km. This result could be partly due to sex differences in the “competition intentions” of the participants: with races < 60 km females reported that they performed the race in a more “pleasure mode” whereas males were more competitively oriented. This possibility should be considered when concluding on fatigability. In other words, less fatigue post-exercise does not necessarily mean a better resistance to fatigue, at least in non-elite runners.

4.2 Muscle Tissue Characteristics

Among the underlying mechanisms, males and females have distinct muscle properties. An obvious difference that mainly explains the difference in maximal strength is muscle mass, due to larger diameter muscle fibers in males rather than the number of fibers [138], this larger size being due to sex-related differences in human skeletal muscle gene (e.g. GRB10 and ACVR2B) expression [139]. This factor has been presented as partly explaining the sex differences in fatigability in most of the studies using single-joint exercise since stronger muscles tend to be less perfused because of higher blood flow occlusion resulting from higher intramuscular pressure. In order to overcome the occlusion effect, Ansdell et al. [140] compared NM fatigue in males and females across submaximal and short duration intermittent isometric contractions (i.e. they used a 3-s contraction duration with a duty cycle allowing to negate the influence of occlusion). After normalizing the intermittent exercise intensity by maximal force, lower force reduction and longer time to task failure was still found in females compared to males. However, this is not the only factor explaining the fatigability. Despite no sex differences in peak force, in power and shortening velocity of single fibers when normalized to muscle cell size (e.g. [141]), there is considerable evidence that females have a greater proportional area of type I fibers in several key muscles for locomotion and daily function [128]. Females also have greater capillarization [100] and lower glycolytic enzyme activity [142] when compared to males. Related to these important features, a better vasodilatory response to exercise has been reported for females. For example, vasodilatory responses of the femoral artery during dynamic knee extensor exercise were found to be greater in

females than in males [143]. Overall, these results explain why females show attenuated impairments in contractile function compared to males [144], and are able to recover at a faster rate from repeated fatiguing exercise [145].

In addition to directly explaining the sex-related difference in fatigability at the muscle level, the attenuated metabolic disturbance related to the greater proportional area of type I may also lead to attenuated afferent feedback (groups III and IV). This may induce less inhibitory inputs to the motoneuronal pool, and thereby less central fatigue. Indeed, greater levels of fatigue observed in males have been associated with greater central deficits, but these studies were using maximal efforts and it is not known if this is also true for submaximal contractions [128].

In conclusion, as generally reported following single-joint fatiguing tasks, females appear less fatigued following trail and ultra-trail races. A greater proportion of type I muscle fibres and higher muscle capillarization (i.e. less muscle de-oxygenation) in females compared to males are possible explanations for lower fatigability in females, and could be of advantage for females over ultra-endurance running performance (Fig. 1). Although the results obtained with the scale on “the competitive intention” of the runners must be treated with caution (i.e. the scale has not undergone a validation process), the fact that females (non-elite) have been showed to be less competition oriented than males could also at least partly explain sex differences in fatigability following such trail running races.

4.3 Muscle Fiber Damage, Inflammation and Delayed-Onset Muscle Soreness

Endurance running races induce metabolic stress and include thousands of ground impacts and subsequent eccentric muscle actions that are likely to induce very focal and limited ultrastructural muscle damage. In the most severe case, less than 10% of the sarcomeres appeared damaged [146], which clearly differ from those induced by pure eccentric muscle actions [147]. The subsequent ultrastructural recovery process involves oedema, remodelling/inflammation, and delayed-onset muscle soreness (DOMS) [148, 149]. The ultrastructural damage is expected to be less or even absent in trained runners [150, 151]. Importantly, (1) neither the degree nor the timing of exercise-induced muscle damage (EIMD) per se correlates well with the DOMS sensation [152, 153], and (2) the timing of DOMS disappearance usually occurs prior to complete structural and functional recoveries [149]. The only study that has directly addressed in humans (via muscle biopsies) the effect of sex on ultrastructural muscle damage and/or inflammation up to 6 days after an eccentric exercise reported no sex difference in muscle damage, yet an attenuated inflammatory response in females as compared to males [154]. In our own study

of the sex differences after a graded 20-km running race [135], DOMS were recorded for the major lower limb muscles. Independently of sex, DOMS was reported at day 2 for the quadriceps muscle group, but only in female runners for the hamstrings, suggesting that possible sex differences are muscle-group dependent. At day 4, despite the disappearance of DOMS, males still had functional deficits in MVC and drop jump test. Female runners also had functional deficits in DJ, but no longer reported DOMS and had already recovered in MVC.

Sorichter et al. [155] have summarized the literature on the identification, usefulness and limits of indirect markers of ultrastructural muscle damage widely used in males and females. Aside from DOMS, the most reported indirect indicators include myoglobin (Mb) and lactate dehydrogenase (LDH) plasma levels as well as serum creatine kinase (CK) activity [156]. According to Goodman et al. [157], the increased serum levels of CK and Mb after an endurance run might result from free radical-induced cell membrane damage and/or transiently increased permeability rather than from mechanical ultrastructural damage. Part of these perturbations are likely to decrease with training (e.g. Margaritis et al. [158]). However, owing to high inter-individual variability none of these indirect markers accurately reflect the magnitude of muscle damage. This appears clearly in the extremely large range of CK concentration values (1500–264,300 IU L⁻¹) reported among finishers of ultra-trail running events, but no differences between sexes have been reported [136, 159]. These methodological limits apply to the serum CK [160], which is still widely used as a marker of the sex difference in the exercise-induced muscle damage [161].

As reviewed by Enns and Tiidus [162], sex and oestrogen are reported as potentially attenuating indices of exercise-induced skeletal muscle damage by influencing the inflammation and repair processes. Although the underlying mechanisms are not yet fully elucidated in humans, oestrogens are thought to exert their protective effects by: (1) acting as an antioxidant, thus limiting oxidative damage; (2) acting as a membrane stabilizer by intercalating within membrane phospholipids; and (3) binding to oestrogen receptors, thus governing downstream muscle repair processes, including the activation and proliferation of muscle satellite cells. Supporting the oestrogen protective effect in runners, lower resting serum CK values are reported in female compared to male athletes [161] and in eumenorrhic compared to amenorrhic female athletes [163], whereas no sex effect was found in the peak values after a marathon [164] in moderately trained runners. Illustrating the limitations of using indirect markers of muscle damage and inflammation, the downhill running study of Oosthuysen and Bosch [165] found that DOMS peaked after 24 h in males and females, but the time to disappearance of CK and DOMS then differed by

sex. Compared with males, females showed earlier restoration of the pre-run CK activity level (at 48 vs. 72 h) but longer-lasting DOMS (up to 72 vs. 48 h).

In conclusion, despite the lack of direct assessment of sex differences in the running-induced ultrastructural muscle damage, the reported time-evolution of indirect markers of muscle inflammation (such as plasmatic CK activity and DOMS) during the recovery period suggest less damage or at least earlier structural recovery in female runners.

4.4 Change in Running Biomechanics and Economy with Fatigue

An altered RE has been repeatedly reported after prolonged road and trail exercises in male runners (e.g. [166–170]), but only a few studies have been conducted in trained or recreational female runners [133]. After a 2-h treadmill run session at a given relative intensity, Glace et al. [133] showed an increase in oxygen cost in male but not in female distance runners. In male athletes, the exacerbated increase in oxygen cost was related to their greater muscle fatigue characterized by the larger strength decrement of the knee extensors (see Sect. 4.1). To the best of our knowledge, only two other studies have examined sex differences with regard to RE change with fatigue, albeit with shorter running bouts. Similar changes in RE were found for males and females following (1) a 1-h run at marathon pace [171] and (2) a 5-km run at 80–85% of $\dot{V}O_{2max}$ [172]. These three experiments were conducted on a treadmill in laboratory conditions. A recent study from our group reported an increase in RE following trail running races from 40 to 170 km; however, no significant sex difference were observed in the changes in RE measured on both flat (+3.1 vs. +8.7% in females vs. males, respectively) and uphill (+1.4 vs. +6.9% in females vs. males, respectively) conditions [137]. Further investigations are warranted to further examine sex differences in RE in the ecological context using models of prolonged road and trail run exercises.

While many studies have investigated sex differences in running biomechanics under acute conditions (see Sect. 2.2), the literature is very limited on sex differences in the alteration of running biomechanics following a fatiguing task. Bazuelo-Ruiz et al. [41] reported a reduced ankle dorsiflexion angle at touchdown in females only after a succession of three consecutive tasks at high intensities (shuttle-run, running up and downstairs, and jumps), this parameter being associated with a decrease in loading rate and vertical impact peak force. These authors suggested a potential adaptative strategy to optimize shock attenuation and thus prevent running injuries. However, these results should be interpreted with caution since the fatiguing protocol was quite unusual. After a 110-km trail running race, Giandolini et al. [173] reported that males adopted a flatter

pattern (i.e. a decreased foot/ground angle) and increased step frequency contrary to females, who landed with a more dorsiflexed ankle accentuating their rear-foot strike pattern. Changes in males were considered as a compensatory adjustment to possibly greater decreased muscle capacity compared to females [173]. Although female adaptations to fatigue observed in the studies of Bazuelo-Ruiz et al. [41] and Giandolini et al. [173] seem to go in opposite directions (concerning the alteration of ankle angle with fatigue), the fatiguing tasks differed considerably between the two studies and could explain the discrepancies in the results. As also suggested by the sex-dependent DOMS sensations per muscle after a graded running race [135] (see Sect. 4.3), it appears that males and females adjust their running pattern differently, which suggests sex-relative biomechanical adaptations. Under laboratory conditions, no sex differences were reported in running biomechanical parameters previously associated with the aetiology or exacerbation of patellofemoral joint pain after a run on a treadmill at 3.5 m s^{-1} until reaching a RPE of 17 out of 20 [68]. These studies compared males and females following different modalities of exercise and under different methodological approaches. Further studies are needed to better investigate potential sex differences in biomechanical alterations following prolonged running exercise.

5 Other Factors Influencing Running Performance

5.1 Thermoregulation

Whether sex differences exist in thermoregulation was uncertain until recently [174], i.e. it was not known if thermoregulatory differences between males and females were due to sex per se or to other differences such as body size or aerobic fitness. Indeed, a correlation between $\dot{V}O_{2\text{max}}$ (lower in females) and the ability to maintain lower core temperatures during exercise in the heat has been found, potentially related to skin blood flow. Females may generate less heat in high ambient temperatures because of lower muscle mass and smaller body size but may be disadvantaged as their lower $\dot{V}O_{2\text{max}}$ can result in faster increases in body temperature. Notley et al. [175] concluded that, contrary to popular belief, sex does not significantly impact exercise thermoregulation [175]. A review by Yanovich et al. [176] confirmed the lack of sex differences and concluded that females were not at greater risk of heat illness, provided the usual risk-management techniques are in place (pacing, clothing, hydration, acclimation). It has been reported that female athletes used more heat illness prevention strategies than male athletes [177]. It is noteworthy that the American College of Sports

Medicine does not provide specific guidelines or strategies for the female athlete. However, it has been argued that a sex difference in thermoregulation may depend on the percentage of humidity and exercise intensity [178]. Indeed, Shapiro et al. [179] found that core temperature increased less in females than in males for hot/humid environments (due to larger surface area to body mass ratio and more efficient sweating suppression), whereas the opposite was true in hot/dry climates (due to lower overall sweating capacity in females despite a higher density of sweat glands). Although thermoregulatory responses vary over the cycle, Notley et al. [175] suggested that there was no additional thermoregulatory concern for female athletes if an oral contraceptive was used or at any phase of the menstrual cycle phase. It is not known if potential sex differences exist in the time-course of acclimation.

It has also been suggested that performance was less affected by warm weather in females than in males during a 161-km ultra-marathon [180], although no sex difference was found in marathons. The opposite may actually be true since performance is more negatively affected for slower populations of runners on that distance [181]. Importantly, because of their smaller body size, lower heat generation, and less sweat loss (i.e. better efficiency in sweating that may decrease the total amount of so-called “wasted sweating” in a humid environment), the water intake required to maintain hydration and thermal balance is attenuated in females. Since hydration guidelines have been developed using data mainly collected in studies with males, they should be interpreted with caution when applied to females [176]. This is particularly important in ultra-endurance exercises since over-hydration may induce gastro-intestinal distress.

5.2 Pacing

Several studies, including a large study with data from 14 US marathons encompassing ~92,000 performances [182], have shown that females tend to have a more even pacing strategy in marathon races [183]. A similar finding has been found in half-marathons [184], 10-km races (at least in fast runners [185]) or even shorter races such as 5-km cross-country race when males and females have similar relative performance [186]. It has been suggested that sex difference in pacing partly reflects a sex difference in decision making [185] rather than greater susceptibility to glycogen depletion [186]. Interestingly, the sex differences in pacing were also found to be true for top runners. A study collected finishing and split times in 673 males and 549 females across nine Olympic and World Championship marathons and found that females slowed less and were more likely to run a negative split than males [187]. In addition, in a study from Trube et al. [188], the more even pacing strategy in females was

found to be an even greater advantage during warm weather conditions.

The number of studies on sex differences in pacing strategy related to ultramarathon is low and there is no clear consistency among the results. Indeed, females showed lower relative starting speeds and higher finishing speeds than males on a 100-km race [189], yet no sex differences in pacing strategy [190] or even better preservation of mean speed throughout the race have been found in males [191] on 24-h races. It has been advised that, given the possibility of a “herd principle”, comparison of competition dynamics between sexes must be further explored in ultra-marathon.

The level of performance and age may be confounding factors in pacing strategies although, again, the results are not that clear. As stated in the present section, sex differences in pacing on 10-k races were mostly found in good runners [185], whereas they occurred across all finishing time groups in the marathon [182]. Deaner et al. [182] also showed that sex differences in pacing occurred across age groups. On the contrary, it has been found that females showed no differences in pace variability across ages, whereas younger (< 30 years) and older (> 60 years) males showed a greater variability in pacing than other age groups [192].

6 Conclusion

Through this narrative review, we have identified a number of sex differences in running related to biomechanical, physiological and NM aspects, some of which potentially play a role in endurance running performance (Fig. 1). In particular, the current literature reports that females show (1) greater hip and knee joint motion in non-sagittal planes and (2) greater lower limb muscle activation, compared to their male counterparts at a given absolute speed. However, most of these sex differences disappear when normalized either by maximal aerobic velocity or anthropometry. Despite those sex differences in running biomechanics, males and females do not differ in terms of RE, suggesting that both sexes optimize their running pattern to adjust to their characteristics. During prolonged exercise, females use more fatty acids than males, which allows them to save carbohydrates. In addition, attenuated NM fatigue, particularly less peripheral impairments on plantar flexor muscles, is observed in females compared to males following endurance and ultra-endurance running bouts. The fact that oestrogen seems to have some protective effects may indicate that females could be less prone to damage induced by repetitive eccentric muscle actions. Yet these differences do not allow females to outrun males in endurance and ultra-endurance running exercises. It is also still not clear whether prolonged running affects males and females differently in terms of running

biomechanics and economy. Sex differences in a fatigued condition are a clearly under-investigated research area and further direct physiological and biomechanical comparisons between males and females following prolonged running are warranted. Considering that females used different pacing (i.e. more even) strategies than males, other psychological and/or sociological parameters that could affect performance or help to understand sex differences in running participation also need to be explored.

Acknowledgements The authors sincerely thank Callum Brownstein for English editing.

Declarations

Author contributions TB, RM, CN, FV and GYM drafted the manuscript; JR, CM and YK provided additional comments and contributions; all authors approved the final version.

Funding No sources of funding were used to assist in the preparation of this article.

Conflict of interest Thibault Besson, Robin Macchi, Jérémy Rossi, Cédric Morio, Yoko Kunimasa, Caroline Nicol, Fabrice Vercauysen and Guillaume Millet declare that they have no conflicts of interest relevant to the content of this review.

References

1. Hafstad AD, Boardman N, Lund J, Hagve M, Wisloff U, Larsen TS, et al. Exercise-induced increase in cardiac efficiency: the impact of intensity. *Circulation*. 2009;120(18):S880.
2. Hespanhol Junior LC, Pillay JD, van Mechelen W, Verhagen E. Meta-analyses of the effects of habitual running on indices of health in physically inactive adults. *Sports Med*. 2015;45(10):1455–68. <https://doi.org/10.1007/s40279-015-0359-y>.
3. van Mechelen W. Running injuries. A review of the epidemiological literature. *Sports Med*. 1992;14(5):320–35.
4. Hanold MT. Beyond the marathon: (De) construction of female ultrarunning bodies. *Sociol Sport J*. 2010;27(2):160–77.
5. Costello JT, Bieuzen F, Bleakley CM. Where are all the female participants in sports and exercise medicine research? *Eur J Sport Sci*. 2014;14(8):847–51.
6. Nuzzo J. Volunteer bias and female participation in exercise and sports science research. *Quest*. 2021;73(1):82–101.
7. Hoffman MD. Performance trends in 161-km ultramarathons. *Int J Sports Med*. 2010;31(1):31–7. <https://doi.org/10.1055/s-0029-1239561>.
8. Whipp BJ, Ward SA. Will women soon outrun men? *Nature*. 1992;355(6355):25. <https://doi.org/10.1038/355025a0>.
9. Tatem AJ, Guerra CA, Atkinson PM, Hay SI. Athletics: momentous sprint at the 2156 Olympics? *Nature*. 2004;431(7008):525. <https://doi.org/10.1038/431525a>.
10. Ochert A. Could women take a lead over men in the long run? *Nature*. 1996;382(6586):15–6. <https://doi.org/10.1038/382015a0>.
11. Bam J, Noakes TD, Juritz J, Dennis SC. Could women outrun men in ultramarathon races? *Med Sci Sports Exerc*.

- 1997;29(2):244–7. <https://doi.org/10.1097/00005768-199702000-00013>.
12. Speechly DP, Taylor SR, Rogers GG. Differences in ultra-endurance exercise in performance-matched male and female runners. *Med Sci Sports Exerc.* 1996;28(3):359–65.
13. Chevront SN, Carter R, Deruisseau KC, Moffatt RJ. Running performance differences between men and women: an update. *Sports Med (Auckland, NZ).* 2005;35(12):1017–24. <https://doi.org/10.2165/00007256-200535120-00002>.
14. Coast JR, Blevins JS, Wilson BA. Do gender differences in running performance disappear with distance? *Can J Appl Physiol.* 2004;29(2):139–45.
15. Lepers R, Cattagni T. Do older athletes reach limits in their performance during marathon running? *Age (Dordr).* 2012;34(3):773–81. <https://doi.org/10.1007/s11357-011-9271-z>.
16. Knechtle B, Rust CA, Rosemann T, Lepers R. Age-related changes in 100-km ultra-marathon running performance. *Age (Dordr).* 2012;34(4):1033–45. <https://doi.org/10.1007/s11357-011-9290-9>.
17. da Fonseca-Engelhardt K, Knechtle B, Rust CA, Knechtle P, Lepers R, Rosemann T. Participation and performance trends in ultra-endurance running races under extreme conditions—'Spartathlon' versus "Badwater." *Extrem Physiol Med.* 2013;2(1):15. <https://doi.org/10.1186/2046-7648-2-15>.
18. Peter L, Rust CA, Knechtle B, Rosemann T, Lepers R. Sex differences in 24-hour ultra-marathon performance—a retrospective data analysis from 1977 to 2012. *Clinics (Sao Paulo).* 2014;69(1):38–46. [https://doi.org/10.6061/clinics/2014\(01\)06](https://doi.org/10.6061/clinics/2014(01)06).
19. Tiller NB, Elliott-Sale KJ, Knechtle B, Wilson PB, Roberts JD, Millet GY. Do sex differences in physiology confer a female advantage in ultra-endurance sport? *Sports Med.* 2021. <https://doi.org/10.1007/s40279-020-01417-2>.
20. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000;32(1):70–84. <https://doi.org/10.1097/00005768-200001000-00012>.
21. McLaughlin JE, Howley ET, Bassett DR, Thompson DL, Fitzhugh EC. Test of the classic model for predicting endurance running performance. *Med Sci Sports Exerc.* 2010;42(5):991–7. <https://doi.org/10.1249/MSS.0b013e3181c0669d>.
22. Davies CT, Thompson MW. Aerobic performance of female marathon and male ultramarathon athletes. *Eur J Appl Physiol Occup Physiol.* 1979;41(4):233–45.
23. Helgerud J. Maximal oxygen uptake, anaerobic threshold and running economy in women and men with similar performances level in marathons. *Eur J Appl Physiol Occup Physiol.* 1994;68(2):155–61. <https://doi.org/10.1007/BF00244029>.
24. Helgerud J, Ingjer F, Stromme SB. Sex differences in performance-matched marathon runners. *Eur J Appl Physiol Occup Physiol.* 1990;61(5–6):433–9.
25. Maughan RJ, Leiper JB. Aerobic capacity and fractional utilisation of aerobic capacity in elite and non-elite male and female marathon runners. *Eur J Appl Physiol Occup Physiol.* 1983;52(1):80–7. <https://doi.org/10.1007/BF00429030>.
26. Sparling PB, Cureton KJ. Biological determinants of the sex difference in 12-min run performance. *Med Sci Sports Exerc.* 1983;15(3):218–23.
27. Calbet JA, Joyner MJ. Disparity in regional and systemic circulatory capacities: do they affect the regulation of the circulation? *Acta Physiol (Oxf).* 2010;199(4):393–406. <https://doi.org/10.1111/j.1748-1716.2010.02125.x>.
28. Carter SL, Rennie C, Tarnopolsky MA. Substrate utilization during endurance exercise in men and women after endurance training. *Am J Physiol Endocrinol Metab.* 2001;280(6):E898–907. <https://doi.org/10.1152/ajpendo.2001.280.6.E898>.
29. Phillips SM, Atkinson SA, Tarnopolsky MA, MacDougall JD. Gender differences in leucine kinetics and nitrogen balance in endurance athletes. *J Appl Physiol (1985).* 1993;75(5):2134–41. <https://doi.org/10.1152/jappl.1993.75.5.2134>.
30. McKenzie S, Phillips SM, Carter SL, Lowther S, Gibala MJ, Tarnopolsky MA. Endurance exercise training attenuates leucine oxidation and BCOAD activation during exercise in humans. *Am J Physiol Endocrinol Metab.* 2000;278(4):E580–7. <https://doi.org/10.1152/ajpendo.2000.278.4.E580>.
31. Cureton KJ. Matching of male and female subjects using VO_{2max} . *Res Q Exerc Sport.* 1981;52(2):264–8. <https://doi.org/10.1080/02701367.1981.10607865>.
32. Sparling PB. A meta-analysis of studies comparing maximal oxygen uptake in men and women. *Res Q Exerc Sport.* 1980;51(3):542–52. <https://doi.org/10.1080/02701367.1980.10608077>.
33. Joyner MJ. Physiological limits to endurance exercise performance: influence of sex. *J Physiol.* 2017;595(9):2949–54.
34. Cureton K, Bishop P, Hutchinson P, Newland H, Vickery S, Zwiren L. Sex difference in maximal oxygen uptake. Effect of equating haemoglobin concentration. *Eur J Appl Physiol Occup Physiol.* 1986;54(6):656–60. <https://doi.org/10.1007/BF00943356>.
35. Garcia-Pinillos F, Jerez-Mayorga D, Latorre-Roman PA, Ramirez-Campillo R, Sanz-Lopez F, Roche-Seruendo LE. How do amateur endurance runners alter spatiotemporal parameters and step variability as running velocity increases? A sex comparison. *J Hum Kinet.* 2020;72:39–49. <https://doi.org/10.2478/hukin-2019-0098>.
36. Nelson RC, Brooks CM, Pike NL. Biomechanical comparison of male and female distance runners. *Ann NY Acad Sci.* 1977;301:793–807. <https://doi.org/10.1111/j.1749-6632.1977.tb38247.x>.
37. Roche-Seruendo L, Latorre-Román P, Soto-Hermoso V, García-Pinillos F. Do sex and body structure influence spatiotemporal step characteristics in endurance runners? *Sci Sports.* 2019;34(6):412.e1–412.e9.
38. Takabayashi T, Edama M, Nakamura M, Nakamura E, Inai T, Kubo M. Gender differences associated with rearfoot, midfoot, and forefoot kinematics during running. *Eur J Sport Sci.* 2017;17(10):1289–96. <https://doi.org/10.1080/17461391.2017.1382578>.
39. Schache AG, Blanch P, Rath D, Wrigley T, Bennell K. Differences between the sexes in the three-dimensional angular rotations of the lumbo-pelvic-hip complex during treadmill running. *J Sports Sci.* 2003;21(2):105–18. <https://doi.org/10.1080/026404031000070859>.
40. Almonroeder TG, Benson LC. Sex differences in lower extremity kinematics and patellofemoral kinetics during running. *J Sports Sci.* 2017;35(16):1575–81. <https://doi.org/10.1080/02640414.2016.1225972>.
41. Bazuelo-Ruiz B, Durá-Gil JV, Palomares N, Medina E, Llana-Belloch S. Effect of fatigue and gender on kinematics and ground reaction forces variables in recreational runners. *PeerJ.* 2018;6:e4489. <https://doi.org/10.7717/peerj.4489>.
42. Chumanov ES, Wall-Scheffler C, Heiderscheid BC. Gender differences in walking and running on level and inclined surfaces. *Clin Biomech (Bristol, Avon).* 2008;23(10):1260–8. <https://doi.org/10.1016/j.clinbiomech.2008.07.011>.
43. Ferber R, Davis IM, Williams DS 3rd. Gender differences in lower extremity mechanics during running. *Clin Biomech (Bristol, Avon).* 2003;18(4):350–7.
44. Gehring D, Mornieux G, Fleischmann J, Gollhofer A. Knee and hip joint biomechanics are gender-specific in runners with high running mileage. *Int J Sports Med.* 2014;35(2):153–8. <https://doi.org/10.1055/s-0033-1343406>.

45. Hannigan JJ, Osternig LR, Chou L-S. Sex-specific relationships between hip strength and hip, pelvis, and trunk kinematics in healthy runners. *J Appl Biomech.* 2018;34(1):76–81.
46. Malinzak RA, Colby SM, Kirkendall DT, Yu B, Garrett WE. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech (Bristol, Avon).* 2001;16(5):438–45. [https://doi.org/10.1016/s0268-0033\(01\)00019-5](https://doi.org/10.1016/s0268-0033(01)00019-5).
47. Nigg BM, Baltich J, Maurer C, Federolf P. Shoe midsole hardness, sex and age effects on lower extremity kinematics during running. *J Biomech.* 2012;45(9):1692–7. <https://doi.org/10.1016/j.jbiomech.2012.03.027>.
48. Phinyomark A, Hettinga BA, Osis ST, Ferber R. Gender and age-related differences in bilateral lower extremity mechanics during treadmill running. *PLoS ONE.* 2014;9(8): e105246. <https://doi.org/10.1371/journal.pone.0105246>.
49. Sakaguchi M, Ogawa H, Shimizu N, Kanehisa H, Yanai T, Kawakami Y. Gender differences in hip and ankle joint kinematics on knee abduction during running. *Eur J Sport Sci.* 2014;14(Suppl 1):S302–9. <https://doi.org/10.1080/17461391.2012.693953>.
50. Sinclair J, Greenhalgh A, Edmundson CJ, Brooks D, Hobbs SJ. Gender differences in the kinetics and kinematics of distance running: implications for footwear design. *Int J Sports Sci Eng.* 2012;6(2):118–28.
51. Willson JD, Petrowitz I, Butler RJ, Kernozek TW. Male and female gluteal muscle activity and lower extremity kinematics during running. *Clin Biomech (Bristol, Avon).* 2012;27(10):1052–7. <https://doi.org/10.1016/j.clinbiomech.2012.08.008>.
52. Hale R. Factors important to women engaged in vigorous physical activity. *Sports medicine.* Philadelphia: WB Saunders; 1984. p. 250–69.
53. Leetun DT, Ireland ML, Willson JD, Ballantyne BT, Davis IM. Core stability measures as risk factors for lower extremity injury in athletes. *Med Sci Sports Exerc.* 2004;36(6):926–34.
54. Hungerford DS, Barry M. Biomechanics of the patellofemoral joint. *Clin Orthop Relat Res.* 1979;144:9–15.
55. Sinclair J, Taylor PJ. Sex differences in tibiofemoral kinematics. *Hum Mov.* 2014;15(2):105–9.
56. von Tschanner V, Goepfert B. Gender dependent EMGs of runners resolved by time/frequency and principal pattern analysis. *J Electromyogr Kinesiol.* 2003;13(3):253–72.
57. DeMont RG, Lephart SM. Effect of sex on preactivation of the gastrocnemius and hamstring muscles. *Br J Sports Med.* 2004;38(2):120–4. <https://doi.org/10.1136/bjism.2002.000195>.
58. Baur H, Hirschmuller A, Cassel M, Muller S, Mayer F. Gender-specific neuromuscular activity of the M. peroneus longus in healthy runners—a descriptive laboratory study. *Clin Biomech Bristol Avon.* 2010;25(9):938–43. <https://doi.org/10.1016/j.clinbiomech.2010.06.009>.
59. Santuz A, Janshen L, Bruell L, Munoz-Martel V, Taborri J, Rossi S, et al. Sex-specific tuning of modular muscle activation patterns for locomotion in young and older adults. *bioRxiv.* 2021.
60. Isherwood J, Wang H, Sterzing T. Running biomechanics and running shoe perception of Chinese men and women. *Footwear Sci.* 2021;13(1):55–67.
61. McGhee DE, Steele JR. Biomechanics of breast support for active women. *Exerc Sport Sci Rev.* 2020;48(3):99–109. <https://doi.org/10.1249/JES.0000000000000221>.
62. Brown N, Scurr J. Do women with smaller breasts perform better in long-distance running? *Eur J Sport Sci.* 2016;16(8):965–71.
63. Keller TS, Weisberger A, Ray J, Hasan S, Shiavi R, Spengler D. Relationship between vertical ground reaction force and speed during walking, slow jogging, and running. *Clin Biomech (Bristol, Avon).* 1996;11(5):253–9.
64. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med.* 2002;36(2):95–101.
65. Glaviano NR, Kew M, Hart JM, Saliba S. Demographic and epidemiological trends in patellofemoral pain. *Int J Sports Phys Ther.* 2015;10(3):281–90.
66. Boling M, Padua D, Marshall S, Guskiewicz K, Pyne S, Beutler A. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. *Scand J Med Sci Sports.* 2010;20(5):725–30. <https://doi.org/10.1111/j.1600-0838.2009.00996.x>.
67. Sinclair J, Selfe J. Sex differences in knee loading in recreational runners. *J Biomech.* 2015;48(10):2171–5. <https://doi.org/10.1016/j.jbiomech.2015.05.016>.
68. Willson JD, Loss JR, Willy RW, Meardon SA. Sex differences in running mechanics and patellofemoral joint kinetics following an exhaustive run. *J Biomech.* 2015;48(15):4155–9. <https://doi.org/10.1016/j.jbiomech.2015.10.021>.
69. Arampatzis A, Monte GD, Karamanidis K, Morey-Klapsing G, Stafiliadis S, Brüggemann G-P. Influence of the muscle-tendon unit's mechanical and morphological properties on running economy. *J Exp Biol.* 2006;209(17):3345–57. <https://doi.org/10.1242/jeb.02340>.
70. Kubo K, Miyazaki D, Shimoju S, Tsunoda N. Relationship between elastic properties of tendon structures and performance in long distance runners. *Eur J Appl Physiol.* 2015;115(8):1725–33.
71. Kubo K, Tabata T, Ikebukuro T, Igarashi K, Yata H, Tsunoda N. Effects of mechanical properties of muscle and tendon on performance in long distance runners. *Eur J Appl Physiol.* 2010;110(3):507–14. <https://doi.org/10.1007/s00421-010-1528-1>.
72. Fletcher JR, MacIntosh BR. Changes in Achilles tendon stiffness and energy cost following a prolonged run in trained distance runners. *PLoS ONE.* 2018;13(8):e0202026. <https://doi.org/10.1371/journal.pone.0202026>.
73. Fletcher JR, Pfister TR, MacIntosh BR. Energy cost of running and Achilles tendon stiffness in man and woman trained runners. *Physiol Rep.* 2013;1(7):e00178. <https://doi.org/10.1002/phy2.178>.
74. Hansen M. Female hormones: do they influence muscle and tendon protein metabolism? *Proc Nutr Soc.* 2018;77(1):32–41.
75. Westh E, Kongsgaard M, Bojsen-Moller J, Aagaard P, Hansen M, Kjaer M, et al. Effect of habitual exercise on the structural and mechanical properties of human tendon, in vivo, in men and women. *Scand J Med Sci Sports.* 2008;18(1):23–30.
76. Sano K, Nicol C, Akiyama M, Kunimasa Y, Oda T, Ito A, et al. Can measures of muscle–tendon interaction improve our understanding of the superiority of Kenyan endurance runners? *Eur J Appl Physiol.* 2015;115(4):849–59.
77. Lee H, Petrofsky JS, Laymon M, Yim J. A greater reduction of anterior cruciate ligament elasticity in women compared to men as a result of delayed onset muscle soreness. *Tohoku J Exp Med.* 2013;231(2):111–5.
78. Shultz SJ, Sander T, Kirk S, Perrin D. Sex differences in knee joint laxity change across the female menstrual cycle. *J Sports Med Phys Fit.* 2005;45(4):594.
79. Balachandar V, Marciniak JL, Wall O, Balachandar C. Effects of the menstrual cycle on lower-limb biomechanics, neuromuscular control, and anterior cruciate ligament injury risk: a systematic review. *Muscles Ligaments Tendons J.* 2017;7(1):136–46. <https://doi.org/10.1138/mltj/2017.7.1.136>.
80. Khowailed IA, Petrofsky J, Lohman E, Daher N, Mohamed O. 17beta-estradiol induced effects on anterior cruciate ligament laxness and neuromuscular activation patterns in female

- runners. *J Womens Health (Larchmt)*. 2015;24(8):670–80. <https://doi.org/10.1089/jwh.2014.5184>.
81. Boisseau N, Duclos M, Guinot M. La femme sportive. *Sciences et pratiques du sport*. 2009.
 82. Isacco L, Boisseau N. Sex hormones and substrate metabolism during endurance exercise. In: Hackney AC, editor. *Sex hormones, exercise and women: scientific and clinical aspects*. Cham: Springer International Publishing; 2017. p. 35–58.
 83. Lundsgaard A-M, Kiens B. Gender differences in skeletal muscle substrate metabolism—molecular mechanisms and insulin sensitivity. *Front Endocrinol*. 2014;5:195. <https://doi.org/10.3389/fendo.2014.00195>.
 84. Tarnopolsky LJ, MacDougall JD, Atkinson SA, Tarnopolsky MA, Sutton JR. Gender differences in substrate for endurance exercise. *J Appl Physiol* (1985). 1990;68(1):302–8. <https://doi.org/10.1152/jappl.1990.68.1.302>.
 85. Tarnopolsky MA. Gender differences in substrate metabolism during endurance exercise. *Can J Appl Physiol*. 2000;25(4):312–27. <https://doi.org/10.1139/h00-024>.
 86. Tarnopolsky MA. Sex differences in exercise metabolism and the role of 17-beta estradiol. *Med Sci Sports Exerc*. 2008;40(4):648–54. <https://doi.org/10.1249/MSS.0b013e31816212ff>.
 87. Venables MC, Achten J, Jeukendrup AE. Determinants of fat oxidation during exercise in healthy men and women: a cross-sectional study. *J Appl Physiol* (1985). 2005;98(1):160–7. <https://doi.org/10.1152/japplphysiol.00662.2003>.
 88. Lamont LS, McCullough AJ, Kalhan SC. Gender differences in leucine, but not lysine, kinetics. *J Appl Physiol* (1985). 2001;91(1):357–62. <https://doi.org/10.1152/jappl.2001.91.1.357>.
 89. Lamont LS, McCullough AJ, Kalhan SC. Gender differences in the regulation of amino acid metabolism. *J Appl Physiol* (1985). 2003;95(3):1259–65. <https://doi.org/10.1152/japplphysiol.01028.2002>.
 90. van Gent RN, Siem D, van Middelkoop M, van Os AG, Bierma-Zeinstra SMA, Koes BW. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med*. 2007;41(8):469–80. <https://doi.org/10.1136/bjism.2006.033548> (discussion 80).
 91. Constantini NW, Dubnov G, Lebrun CM. The menstrual cycle and sport performance. *Clin Sports Med*. 2005;24(2):e51–82. <https://doi.org/10.1016/j.csm.2005.01.003> (xiii–xiv).
 92. D'Eon TM, Sharoff C, Chipkin SR, Grow D, Ruby BC, Braun B. Regulation of exercise carbohydrate metabolism by estrogen and progesterone in women. *Am J Physiol Endocrinol Metab*. 2002;283(5):E1046–55. <https://doi.org/10.1152/ajpendo.00271.2002>.
 93. Ruby BC, Robergs RA, Waters DL, Burge M, Mermier C, Stolarczyk L. Effects of estradiol on substrate turnover during exercise in amenorrheic females. *Med Sci Sports Exerc*. 1997;29(9):1160–9. <https://doi.org/10.1097/00005768-199709000-00007>.
 94. Hackney AC, McCracken-Compton MA, Ainsworth B. Substrate responses to submaximal exercise in the midfollicular and midluteal phases of the menstrual cycle. *Int J Sport Nutr Exerc Metab*. 1994;4(3):299–308. <https://doi.org/10.1123/ijns.4.3.299>.
 95. Hackney AC, Muoio D, Meyer WR. The Effect of sex steroid hormones on substrate oxidation during prolonged submaximal exercise in women. *Jpn J Physiol*. 2000;50(5):489–94. <https://doi.org/10.2170/jjphysiol.50.489>.
 96. Kanaley JA, Boileau RA, Bahr JA, Misner JE, Nelson RA. Substrate oxidation and GH responses to exercise are independent of menstrual phase and status. *Med Sci Sports Exerc*. 1992;24(8):873–80.
 97. Tarnopolsky MA, Rennie CD, Robertshaw HA, Fedak-Tarnopolsky SN, Devries MC, Hamadeh MJ. Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial ultrastructure, substrate use, and mitochondrial enzyme activity. *Am J Physiol Regul Integr Comp Physiol*. 2007;292(3):R1271–8. <https://doi.org/10.1152/ajpregu.00472.2006>.
 98. Fisher G, Windham ST, Griffin P, Warren JL, Gower BA, Hunter GR. Associations of human skeletal muscle fiber type and insulin sensitivity, blood lipids, and vascular hemodynamics in a cohort of premenopausal women. *Eur J Appl Physiol*. 2017;117(7):1413–22. <https://doi.org/10.1007/s00421-017-3634-9>.
 99. Essén B, Jansson E, Henriksson J, Taylor AW, Saltin B. Metabolic characteristics of fibre types in human skeletal muscle. *Acta Physiol Scand*. 1975;95(2):153–65. <https://doi.org/10.1111/j.1748-1716.1975.tb10038.x>.
 100. Roepstorff C, Donsmark M, Thiele M, Vistisen B, Stewart G, Vissing K, et al. Sex differences in hormone-sensitive lipase expression, activity, and phosphorylation in skeletal muscle at rest and during exercise. *Am J Physiol Endocrinol Metab*. 2006;291(5):E1106–14. <https://doi.org/10.1152/ajpendo.00097.2006>.
 101. Roepstorff C, Steffensen CH, Madsen M, Stallknecht B, Kanstrup I-L, Richter EA, et al. Gender differences in substrate utilization during submaximal exercise in endurance-trained subjects. *Am J Physiol Endocrinol Metab*. 2002;282(2):E435–47. <https://doi.org/10.1152/ajpendo.00266.2001>.
 102. Steffensen CH, Roepstorff C, Madsen M, Kiens B. Myocellular triacylglycerol breakdown in females but not in males during exercise. *Am J Physiol Endocrinol Metab*. 2002;282(3):E634–42. <https://doi.org/10.1152/ajpendo.00078.2001>.
 103. Fletcher JR, MacIntosh BR. Running economy from a muscle energetics perspective. *Front Physiol*. 2017;2017:8. <https://doi.org/10.3389/fphys.2017.00433>.
 104. Moore IS. Is there an economical running technique? A review of modifiable biomechanical factors affecting running economy. *Sports Med*. 2016;46(6):793–807. <https://doi.org/10.1007/s40279-016-0474-4>.
 105. Fletcher JR, Esau SP, MacIntosh BR. Economy of running: beyond the measurement of oxygen uptake. *J Appl Physiol* (1985). 2009;107(6):1918–22. <https://doi.org/10.1152/japplphysiol.00307.2009>.
 106. Vernillo G, Millet GP, Millet GY. Does the running economy really increase after ultra-marathons? *Front Physiol*. 2017. <https://doi.org/10.3389/fphys.2017.00783>.
 107. Romijn JA, Coyle EF, Sidossis LS, Rosenblatt J, Wolfe RR. Substrate metabolism during different exercise intensities in endurance-trained women. *J Appl Physiol* (1985). 2000;88(5):1707–14. <https://doi.org/10.1152/jappl.2000.88.5.1707>.
 108. Barnes KR, Mcguigan MR, Kilding AE. Lower-body determinants of running economy in male and female distance runners. *J Strength Cond Res*. 2014;28(5):1289–97. <https://doi.org/10.1519/JSC.0000000000000267>.
 109. Black MI, Handsaker JC, Allen SJ, Forrester SE, Folland JP. Is there an optimal speed for economical running? *Int J Sports Physiol Perform*. 2018;13(1):75–81. <https://doi.org/10.1123/ijssp.2017-0015>.
 110. Bransford DR, Howley ET. Oxygen cost of running in trained and untrained men and women. *Med Sci Sports Exerc*. 1977;9(1):41–4.
 111. Bunc V, Heller J. Energy cost of running in similarly trained men and women. *Eur J Appl Physiol*. 1989;59(3):178–83. <https://doi.org/10.1007/BF02386184>.
 112. Daniels J, Daniels N. Running economy of elite male and elite female runners. *Med Sci Sports Exerc*. 1992;24(4):483–9.
 113. Fletcher JR, MacIntosh BR. Achilles tendon strain energy in distance running: consider the muscle energy cost. *J Appl Physiol*

- (1985). 2014;118(2):193–9. <https://doi.org/10.1152/jappphysiol.00732.2014>.
114. Morgan DW, Craib M. Physiological aspects of running economy. *Med Sci Sports Exerc.* 1992;24(4):456–61.
 115. Helgerud J, Støren Ø, Hoff J. Are there differences in running economy at different velocities for well-trained distance runners? *Eur J Appl Physiol.* 2010;108(6):1099–105. <https://doi.org/10.1007/s00421-009-1218-z>.
 116. Devries MC, Hamadeh MJ, Phillips SM, Tarnopolsky MA. Menstrual cycle phase and sex influence muscle glycogen utilization and glucose turnover during moderate-intensity endurance exercise. *Am J Physiol Regul Integr Comp Physiol.* 2006;291(4):R1120–8. <https://doi.org/10.1152/ajpregu.00700.2005>.
 117. Mittendorfer B, Horowitz JF, Klein S. Gender differences in lipid and glucose kinetics during short-term fasting. *Am J Physiol Regul Integr Comp Physiol.* 2001;281(6):E1333–9. <https://doi.org/10.1152/ajpendo.2001.281.6.E1333>.
 118. Giacomoni M, Falgairette G. Decreased submaximal oxygen uptake during short duration oral contraceptive use: a randomized cross-over trial in premenopausal women. *Ergonomics.* 2000;43(10):1559–70. <https://doi.org/10.1080/001401300750003989>.
 119. Dokumaci B, Hazir T. Effects of the menstrual cycle on running economy: oxygen cost versus caloric cost. *Res Q Exercise Sport.* 2019;90(3):318–26. <https://doi.org/10.1080/02701367.2019.1599800>.
 120. Goldsmith E, Glaister M. The effect of the menstrual cycle on running economy. *J Sports Med Phys Fit.* 2020;60(4):610–7. <https://doi.org/10.23736/S0022-4707.20.10229-9>.
 121. Williams TJ, Krahenbuhl GS. Menstrual cycle phase and running economy. *Med Sci Sports Exerc.* 1997;29(12):1609–18. <https://doi.org/10.1097/00005768-199712000-00010>.
 122. Barry BK, Enoka RM. The neurobiology of muscle fatigue: 15 years later. *Integr Comp Biol.* 2007;47(4):465–73.
 123. Giandolini M, Vernillo G, Samozino P, Horvais N, Edwards WB, Morin JB, et al. Fatigue associated with prolonged graded running. *Eur J Appl Physiol.* 2016;116(10):1859–73. <https://doi.org/10.1007/s00421-016-3437-4>.
 124. Millet GY. Can neuromuscular fatigue explain running strategies and performance in ultra-marathons? *Sports Med.* 2011;41(6):489–506. <https://doi.org/10.2165/11588760-00000000-00000>.
 125. Noakes TD. The central governor model of exercise regulation applied to the marathon. *Sports Med.* 2007;37(4–5):374–7. <https://doi.org/10.2165/00007256-200737040-00026>.
 126. Thomas K, Elmeua M, Howatson G, Goodall S. Intensity-dependent contribution of neuromuscular fatigue after constant-load cycling. *Med Sci Sports Exerc.* 2016;48(9):1751–60. <https://doi.org/10.1249/MSS.0000000000000950>.
 127. Bontemps B, Vercruyssen F, Gruet M, Louis J. Downhill running: what are the effects and how can we adapt? A narrative review. *Sports Med.* 2020;50(12):2083–110. <https://doi.org/10.1007/s40279-020-01355-z>.
 128. Hunter SK. Sex differences in human fatigability: mechanisms and insight to physiological responses. *Acta Physiol.* 2014;210(4):768–89. <https://doi.org/10.1111/apha.12234>.
 129. Hunter SK. The relevance of sex differences in performance fatigability. *Med Sci Sports Exerc.* 2016;48(11):2247–56. <https://doi.org/10.1249/MSS.0000000000000928>.
 130. Pincivero DM, Gandaio CM, Ito Y. Gender-specific knee extensor torque, flexor torque, and muscle fatigue responses during maximal effort contractions. *Eur J Appl Physiol.* 2003;89(2):134–41. <https://doi.org/10.1007/s00421-002-0739-5>.
 131. Ehrstrom S, Tartaruga MP, Easthope CS, Brisswalter J, Morin JB, Vercruyssen F. Short trail running race: beyond the classic model for endurance running performance. *Med Sci Sports Exerc.* 2018;50(3):580–8. <https://doi.org/10.1249/MSS.000000000001467>.
 132. Ansdell P, Brownstein CG, Skarabot J, Hicks KM, Howatson G, Thomas K, et al. Sex differences in fatigability and recovery relative to the intensity–duration relationship. *J Physiol.* 2019;597(23):5577–95. <https://doi.org/10.1113/JP278699>.
 133. Glace BW, McHugh MP, Gleim GW. Effects of a 2-hour run on metabolic economy and lower extremity strength in men and women. *J Orthop Sports Phys Ther.* 1998;27(3):189–96. <https://doi.org/10.2519/jospt.1998.27.3.189>.
 134. Boccia G, Dardanello D, Tarperi C, Festa L, La Torre A, Pellegrini B, et al. Women show similar central and peripheral fatigue to men after half-marathon. *Eur J Sport Sci.* 2018. <https://doi.org/10.1080/17461391.2018.1442500>.
 135. Macchi R, Vercruyssen F, Hays A, Aubert G, Exubis G, Chavet P, et al. Sex influence on the functional recovery pattern after a graded running race: original analysis to identify the recovery profiles. *Front Physiol.* 2021;12:311.
 136. Temesi J, Arnal PJ, Rupp T, Feasson L, Cartier R, Gergele L, et al. Are females more resistant to extreme neuromuscular fatigue? *Med Sci Sports Exerc.* 2015;47(7):1372–82. <https://doi.org/10.1249/MSS.0000000000000540>.
 137. Besson T, Parent A, Brownstein CG, Espeit L, Lapole T, Martin V, et al. Sex differences in neuromuscular fatigue and changes in cost of running after mountain trail races of various distances. *Med Sci Sports Exerc.* 2021;53(11):2374–87. <https://doi.org/10.1249/MSS.0000000000002719>.
 138. Miller AE, MacDougall JD, Tarnopolsky MA, Sale DG. Gender differences in strength and muscle fiber characteristics. *Eur J Appl Physiol Occup Physiol.* 1993;66(3):254–62. <https://doi.org/10.1007/BF00235103>.
 139. Welle S, Tawil R, Thornton CA. Sex-related differences in gene expression in human skeletal muscle. *PLoS ONE.* 2008;3(1):e1385. <https://doi.org/10.1371/journal.pone.0001385>.
 140. Ansdell P, Thomas K, Howatson G, Hunter S, Goodall S. Contraction intensity and sex differences in knee-extensor fatigability. *J Electromyogr Kinesiol.* 2017;37:68–74. <https://doi.org/10.1016/j.jelekin.2017.09.003>.
 141. Trappe S, Gallagher P, Harber M, Carrithers J, Fluckey J, Trappe T. Single muscle fibre contractile properties in young and old men and women. *J Physiol.* 2003;552(Pt 1):47–58. <https://doi.org/10.1113/jphysiol.2003.044966>.
 142. Komi PV, Bosco C. Utilization of stored elastic energy in leg extensor muscles by men and women. *Med Sci Sports Exerc.* 1978;10(4):261–5.
 143. Parker BA, Smithmyer SL, Pelberg JA, Mishkin AD, Herr MD, Proctor DN. Sex differences in leg vasodilation during graded knee extensor exercise in young adults. *J Appl Physiol (1985).* 2007;103(5):1583–91. <https://doi.org/10.1152/jappphysiol.00662.2007>.
 144. Sundberg CW, Hunter SK, Bundle MW. Rates of performance loss and neuromuscular activity in men and women during cycling: evidence for a common metabolic basis of muscle fatigue. *J Appl Physiol (1985).* 2017;122(1):130–41. <https://doi.org/10.1152/jappphysiol.00468.2016>.
 145. Laurent C, Green J, Bishop P, Sjökvist J, Schumacker R, Richardson M, et al. Effect of gender on fatigue and recovery following maximal intensity repeated sprint performance. *J Sports Med Phys Fit.* 2010;50(3):243–53.
 146. Warhol MJ, Siegel AJ, Evans WJ, Silverman LM. Skeletal muscle injury and repair in marathon runners after competition. *Am J Clin Pathol.* 1985;118(2):331–9.
 147. Paulsen G, Mikkelsen UR, Raastad T, Peake JM. Leucocytes, cytokines and satellite cells: what role do they play in muscle

- damage and regeneration following eccentric exercise? *Exerc Immunol Rev.* 2012;2012(18):42–97.
148. Nicol C, Avela J, Komi PV. The stretch-shortening cycle: a model to study naturally occurring neuromuscular fatigue. *Sports Med.* 2006;36(11):977–99. <https://doi.org/10.2165/00007256-200636110-00004>.
 149. Nicol C, Komi PV. Stretch-shortening cycle fatigue. Neuromuscular aspects of sport performance. New York: Wiley; 2010. p. 183–215.
 150. Evans WJ, Meredith CN, Cannon JG, Dinarello CA, Frontera WR, Hughes VA, et al. Metabolic changes following eccentric exercise in trained and untrained men. *J Appl Physiol* (1985). 1986;61(5):1864–8. <https://doi.org/10.1152/jappl.1986.61.5.1864>.
 151. Siegel A, Silerman L, Lopez R. Creatine kinase elevations in marathon runners: relationship to training and competition. *Yale J Biol Med.* 1980;53(4):275–9.
 152. Newham DJ, Mills KR, Quigley BM, Edwards RH. Pain and fatigue after concentric and eccentric muscle contractions. *Clin Sci (Lond).* 1983;64(1):55–62.
 153. Howell JN, Chleboun G, Conatser R. Muscle stiffness, strength loss, swelling and soreness following exercise-induced injury in humans. *J Physiol.* 1993;464:183–96.
 154. Stupka N, Lowther S, Chorneyko K, Bourgeois J, Hogben C, Tarnopolsky M. Gender differences in muscle inflammation after eccentric exercise. *J Appl Physiol* (1985). 2000;89(6):2325–32.
 155. Sorichter S, Puschendorf B, Mair J. Skeletal muscle injury induced by eccentric muscle action: muscle proteins as markers of muscle fiber injury. *Exerc Immunol Rev.* 1999;1999(5):5–21.
 156. Silva RP, Vilaça A, Guerra FD, Mundim AV, De Agostini GG, De Abreu LC, et al. Sex differences in physiological stress induced by a long-lasting adventure race: a prospective observational analytical study. *Sportverletz Sportschaden.* 2020;34(02):84–95.
 157. Goodman C, Henry G, Dawson B, Gillam I, Beilby J, Ching S, et al. Biochemical and ultrastructural indices of muscle damage after a twenty-one kilometre run. *Aust J Sci Med Sport.* 1997;29(4):95–8.
 158. Margaritis I, Tessier F, Richard M-J, Marconnet P. No evidence of oxidative stress after a triathlon race in highly trained competitors. *Int J Sports Med.* 1997;18(3):186–90.
 159. Hoffman MD, Ingwerson JL, Rogers IR, Hew-Butler T, Stuenkel KJ. Increasing creatine kinase concentrations at the 161-km Western States Endurance Run. *Wilderness Environ Med.* 2012;23(1):56–60. <https://doi.org/10.1016/j.wem.2011.11.001>.
 160. Fridén J, Lieber RL. Serum creatine kinase level is a poor predictor of muscle function after injury. *Scand J Med Sci Sports.* 2001;11(2):126–7. <https://doi.org/10.1034/j.1600-0838.2001.011002126.x>.
 161. Mougios V. Reference intervals for serum creatine kinase in athletes. *Br J Sports Med.* 2007;41(10):674–8. <https://doi.org/10.1136/bjism.2006.034041>.
 162. Enns DL, Tiidus PM. The influence of estrogen on skeletal muscle: sex matters. *Sports Med.* 2010;40(1):41–58. <https://doi.org/10.2165/11319760-000000000-00000>.
 163. Stachenfeld NS, Taylor HS. Challenges and methodology for testing young healthy women in physiological studies. *Am J Physiol Endocrinol Metab.* 2014;306(8):E849–53. <https://doi.org/10.1152/ajpendo.00038.2014>.
 164. Kaikkonen J, Porkkala-Sarataho E, Tuomainen TP, Nyssönen K, Kosonen L, Ristonmaa U, et al. Exhaustive exercise increases plasma/serum total oxidation resistance in moderately trained men and women, whereas their VLDL + LDL lipoprotein fraction is more susceptible to oxidation. *Scand J Clin Lab Invest.* 2002;62(8):599–607. <https://doi.org/10.1080/003655102764654330>.
 165. Oosthuysen T, Bosch AN. The effect of gender and menstrual phase on serum creatine kinase activity and muscle soreness following downhill running. *Antioxidants.* 2017. <https://doi.org/10.3390/antiox6010016>.
 166. Brueckner JC, Atchou G, Capelli C, Duvallet A, Barrault D, Jousset E, et al. The energy cost of running increases with the distance covered. *Eur J Appl Physiol.* 1991;62(6):385–9. <https://doi.org/10.1007/BF00626607>.
 167. Hunter I, Smith GA. Preferred and optimal stride frequency, stiffness and economy: changes with fatigue during a 1-h high-intensity run. *Eur J Appl Physiol.* 2007;100(6):653–61. <https://doi.org/10.1007/s00421-007-0456-1>.
 168. Vercruyssen F, Gruet M, Colson SS, Ehrstrom S, Brisswalter J. Compression garments, muscle contractile function, and economy in trail runners. *Int J Sports Physiol Perform.* 2017;12(1):62–8. <https://doi.org/10.1123/ijsspp.2016-0035>.
 169. Vercruyssen F, Tartaruga M, Horvais N, Brisswalter J. Effects of footwear and fatigue on running economy and biomechanics in trail runners. *Med Sci Sports Exerc.* 2016;48(10):1976–84. <https://doi.org/10.1249/MSS.0000000000000981>.
 170. Xu F, Montgomery DL. Effect of prolonged exercise at 65 and 80 % of VO_{2max} on running economy. *Int J Sports Med.* 1995;16(5):309–13. <https://doi.org/10.1055/s-2007-973011>.
 171. Loftin M, Sothorn M, Tuuri G, Tompkins C, Koss C, Bonis M. Gender comparison of physiologic and perceptual responses in recreational marathon runners. *Int J Sports Physiol Perform.* 2009;4(3):307–16.
 172. Thomas DQ, Fernhall B, Granat H. Changes in running economy during a 5-km run in trained men and women runners. *J Strength Cond Res.* 1999;13(2):162–7.
 173. Giandolini M, Gimenez P, Millet GY, Morin J-B, Samozino P. Consequences of an ultra-trail on impact and lower limb kinematics in male and female runners. *Footwear Sci.* 2013;5(sup1):S14–5.
 174. Notley SR, Racinais S, Kenny GP. Do sex differences in thermoregulation pose a concern for female athletes preparing for the Tokyo Olympics? *Br J Sports Med.* 2021;55(6):298–9. <https://doi.org/10.1136/bjsports-2020-102911>.
 175. Notley SR, Akerman AP, Meade RD, McGarr GW, Kenny GP. Exercise thermoregulation in prepubertal children: a brief methodological review. *Med Sci Sports Exerc.* 2020. <https://doi.org/10.1249/MSS.0000000000002391>.
 176. Yanovich R, Ketko I, Charkoudian N. Sex differences in human thermoregulation: relevance for 2020 and beyond. *Physiology (Bethesda).* 2020;35(3):177–84. <https://doi.org/10.1152/physiol.00035.2019>.
 177. Periard JD, Racinais S, Timpka T, Dahlstrom O, Spreco A, Jacobsson J, et al. Strategies and factors associated with preparing for competing in the heat: a cohort study at the 2015 IAAF World Athletics Championships. *Br J Sports Med.* 2017;51(4):264–70. <https://doi.org/10.1136/bjsports-2016-096579>.
 178. Gagnon D, Crandall CG, Kenny GP. Sex differences in post-synaptic sweating and cutaneous vasodilation. *J Appl Physiol* (1985). 2013;114(3):394–401. <https://doi.org/10.1152/japplphysiol.00877.2012>.
 179. Shapiro Y, Pandolf KB, Avellini BA, Pimental NA, Goldman RF. Physiological responses of men and women to humid and dry heat. *J Appl Physiol Respir Environ Exerc Physiol.* 1980;49(1):1–8. <https://doi.org/10.1152/jappl.1980.49.1.1>.
 180. Wegelin JA, Hoffman MD. Variables associated with odds of finishing and finish time in a 161-km ultramarathon. *Eur J Appl Physiol.* 2011;111(1):145–53. <https://doi.org/10.1007/s00421-010-1633-1>.

181. Ely MR, Cheuvront SN, Roberts WO, Montain SJ. Impact of weather on marathon-running performance. *Med Sci Sports Exerc.* 2007;39(3):487–93. <https://doi.org/10.1249/mss.0b013e31802d3aba>.
182. Deaner RO, Carter RE, Joyner MJ, Hunter SK. Men are more likely than women to slow in the marathon. *Med Sci Sports Exerc.* 2015;47(3):607–16. <https://doi.org/10.1249/MSS.000000000000432>.
183. March DS, Vanderburgh PM, Titlebaum PJ, Hoops ML. Age, sex, and finish time as determinants of pacing in the marathon. *J Strength Cond Res.* 2011;25(2):386–91. <https://doi.org/10.1519/JSC.0b013e3181bffd0f>.
184. Cuk I, Nikolaidis PT, Knechtle B. Sex differences in pacing during half-marathon and marathon race. *Res Sports Med.* 2020;28(1):111–20. <https://doi.org/10.1080/15438627.2019.1593835>.
185. Deaner RO, Addona V, Carter RE, Joyner MJ, Hunter SK. Fast men slow more than fast women in a 10 kilometer road race. *PeerJ.* 2016;4:e2235. <https://doi.org/10.7717/peerj.2235>.
186. Deaner RO, Lowen A. Males and females pace differently in high school cross-country races. *J Strength Cond Res.* 2016;30(11):2991–7. <https://doi.org/10.1519/JSC.0000000000001407>.
187. Hanley B. Pacing, packing and sex-based differences in Olympic and IAAF World Championship marathons. *J Sports Sci.* 2016;34(17):1675–81. <https://doi.org/10.1080/02640414.2015.1132841>.
188. Trubee NW, Vanderburgh PM, Diestelkamp WS, Jackson KJ. Effects of heat stress and sex on pacing in marathon runners. *J Strength Cond Res.* 2014;28(6):1673–8. <https://doi.org/10.1519/JSC.000000000000295>.
189. Renfree A, Crivoi do Carmo E, Martin L. The influence of performance level, age and gender on pacing strategy during a 100-km ultramarathon. *Eur J Sport Sci.* 2016;16(4):409–15. <https://doi.org/10.1080/17461391.2015.1041061>.
190. Bossi AH, Matta GG, Millet GY, Lima P, Pertence LC, de Lima JP, et al. Pacing strategy during 24-hour ultramarathon-distance running. *Int J Sports Physiol Perform.* 2017;12(5):590–6. <https://doi.org/10.1123/ijsp.2016-0237>.
191. Inoue A, Santos TM, Hettinga FJ, de Souza Alves D, Viana BF, de Souza Terra B, et al. The impact of sex and performance level on pacing behavior in a 24-h ultramarathon. In: *Hurdling the Challenges of the 2019 IAAF World Championships.* 2020
192. Cuk I, Nikolaidis PT, Markovic S, Knechtle B. Age differences in pacing in endurance running: comparison between marathon and half-marathon men and women. *Medicina (Kaunas).* 2019. <https://doi.org/10.3390/medicina55080479>.
193. Esculier JF, Willy RW, Baggaley MW, Meardon SA, Willson JD. Sex-specific kinetic and kinematic indicators of medial tibiofemoral force during walking and running. *Knee.* 2017;24(6):1317–25. <https://doi.org/10.1016/j.knee.2017.08.054>.
194. Hennig E. Gender differences for running in athletic footwear. In: *Proceedings of the 5th Symposium on Footwear biomechanics, Zurich, Switzerland; 2001.* p. 44–5.