



**HAL**  
open science

## Metabolic challenges and key players in serpentinite-hosted microbial ecosystems

Rabja Maria Popall, Anne Postec, Aurélien Lecoivre, Marianne Quéméneur,  
Gaël Erauso

► **To cite this version:**

Rabja Maria Popall, Anne Postec, Aurélien Lecoivre, Marianne Quéméneur, Gaël Erauso. Metabolic challenges and key players in serpentinite-hosted microbial ecosystems. *Frontiers in Microbiology*, 2023, 14, 10.3389/fmicb.2023.1197823 . hal-04172274

**HAL Id: hal-04172274**

**<https://amu.hal.science/hal-04172274v1>**

Submitted on 12 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## OPEN ACCESS

## EDITED BY

William J. Brazelton,  
The University of Utah, United States

## REVIEWED BY

Daniel Colman,  
Montana State University, United States  
Leah Trutschel,  
University of Cincinnati, United States

## \*CORRESPONDENCE

Gaël Erauso  
✉ gael.erauso@mio.osupytheas.fr

RECEIVED 31 March 2023

ACCEPTED 29 June 2023

PUBLISHED 24 July 2023

## CITATION

Popall RM, Postec A, Lecoivre A,  
Quéméneur M and Erauso G (2023) Metabolic  
challenges and key players in serpentinite-  
hosted microbial ecosystems.  
*Front. Microbiol.* 14:1197823.  
doi: 10.3389/fmicb.2023.1197823

## COPYRIGHT

© 2023 Popall, Postec, Lecoivre, Quéméneur  
and Erauso. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in this  
journal is cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Metabolic challenges and key players in serpentinite-hosted microbial ecosystems

Rabja Maria Popall, Anne Postec, Aurélien Lecoivre,  
Marianne Quéméneur and Gaël Erauso\*

Aix-Marseille Univ, Univ Toulon, CNRS, IRD, MIO, Marseille, France

Serpentinite-hosted systems are amongst the most challenging environments for life on Earth. Serpentinization, a geochemical alteration of exposed ultramafic rock, produces hydrothermal fluids enriched in abiotically derived hydrogen (H<sub>2</sub>), methane (CH<sub>4</sub>), and small organic molecules. The hyperalkaline pH of these fluids poses a great challenge for metabolic energy and nutrient acquisition, curbing the cellular membrane potential and limiting electron acceptor, carbon, and phosphorous availability. Nevertheless, serpentinization supports the growth of diverse microbial communities whose metabolic make-up might shed light on the beginning of life on Earth and potentially elsewhere. Here, we outline current hypotheses on metabolic energy production, carbon fixation, and nutrient acquisition in serpentinizing environments. A taxonomic survey is performed for each important metabolic function, highlighting potential key players such as H<sub>2</sub> and CH<sub>4</sub> cycling *Serpentinimonas*, *Hydrogenophaga*, *Methanobacteriales*, *Methanosarcinales*, and novel candidate phyla. Methodological biases of the available data and future approaches are discussed.

## KEYWORDS


serpentinization, alkaline hydrothermal system, alkaliphile, hydrogenotroph, lithotroph, submarine alkaline vent theory, origin of life

## 1. Introduction



The beginnings of life remain one of the most outstanding scientific issues and have been dubbed the “black hole at the heart of biology” (Lane, 2015). One of the central requirements for living systems is a continuous physicochemical disequilibrium driving biological activity (Russell et al., 1988). In marine alkaline hydrothermal systems, strong electrochemical gradients develop between the ultrabasic hydrothermal fluid rising from the deep subsurface, and the seawater. These gradients are maintained across the porous hydrothermal chimney wall, which can be compared to an osmotic membrane. It is hypothesized that this rudimentary proton motive force has driven chimney nanopores to develop into protocells at the emergence of life (Russell et al., 2010; Sojo et al., 2016). Modern ecosystems at hydrothermal vents might thus provide a glimpse into very early microbial life forms.

Most alkaline hydrothermal systems are formed in environments where mantle rocks have been tectonically uplifted and exposed, either above sea level or on the seafloor. Contact with water initiates serpentinization, a geochemical alteration of the ultramafic rock. This process

yields large amounts of hydrogen (H<sub>2</sub>) and constitutes one of the most important sources of H<sub>2</sub> on Earth (Reaction 1) (Truche et al., 2020).



The oxidation of ferrous iron in olivine or pyroxene to ferric iron and magnetite by water creates reducing conditions. Catalyzed by minerals, this facilitates abiotic reactions of the produced H<sub>2</sub> with mantle-derived carbon dioxide (CO<sub>2</sub>) or carbon monoxide (CO) (McCollom and Seewald, 2001). In Sabatier (Reaction 2) and Fischer-Tropsch (Reaction 3) type processes, methane (CH<sub>4</sub>) and small organic molecules [C<sub>n</sub>H(2n+2)] are enriched in the hydrothermal fluid (Barbier et al., 2020):




The serpentinization reaction produces hydrothermal fluids with pH values commonly surpassing 12. In these ultrabasic conditions, the carbonate equilibrium is permanently shifted from CO<sub>2</sub> to carbonate species, removing most dissolved inorganic carbon (DIC) from the environment. Upon reaching the rock surface, much of the carbonate precipitates with fluid-derived calcium. Over time, calcium carbonate amalgamates with brucite and forms the chimneys or travertines typical for serpentinizing environments (Barnes et al., 1978; Früh-Green et al., 2004; McCollom and Seewald, 2013) (Figure 1A).

## 2. Serpentinite-hosted ecosystems

The products of serpentinization can support chemosynthetic microbial ecosystems growing independently from sunlight. Such serpentinite-hosted ecosystems are found in marine and continental environments, with hydrothermal fluids originating from marine, meteoric and/or groundwater sources.

Most marine serpentinizing ecosystems are located along rather slow-spreading mid-ocean ridges, where continuous tectonic activity facilitates frequent exposure of ultramafic rock (Schrenk et al., 2013; Albers et al., 2021). The most prominent example is the Lost City hydrothermal field near the Mid-Atlantic ridge (e.g., Kelley et al., 2005). A similar system, the Old City hydrothermal field, has recently been discovered along the Southwest Indian ridge (Lecoivre et al., 2021). Several other marine sites host mixed-type ecosystems that feature characteristics of both alkaline and acidic hydrothermal vents, such as the Rainbow (Flores et al., 2011), Logatchev (Perner et al., 2007), Ashadze (Fabri et al., 2011) and Kairei fields (Nakamura et al., 2009).

The most well-studied continental systems include the Samail ophiolite in Oman (Rempfert et al., 2017; Kraus et al., 2021), the Tablelands (Bay of Islands) ophiolite in Newfoundland (Brazelton et al., 2013), the Cabeço de Vide aquifer in Portugal (Tiago and Verissimo, 2013), the Leka ophiolite complex in Norway (Daae et al., 2013), the Italian Gruppo di Voltri (Quéméneur et al., 2015), the Zambales ophiolite in the Philippines (Woycheese et al., 2015), The Cedars (Suzuki et al., 2013; Kohl et al., 2016) and Coast Range (Twing et al., 2017) ophiolites on the West Coast of the United States, the Chimaera (Tekirova) ophiolite in Turkey (Neubeck et al., 2017), the tropical Santa Elena ophiolite in Costa Rica (Crespo-Medina et al., 2017), the Del Puerto ophiolite in California (Blank et al., 2009) and the Troodos ophiolite in Cyprus (Rizoulis et al., 2016).



**FIGURE 1**  
The biogeochemical environment of marine serpentinite-hosted systems. **(A)** Production of compounds of interest in microbial metabolism. Reactions are color-coded with solid arrows showing abiotic processes, empty arrows showing biotic processes, and striped arrows indicating external supply. The metabolic potential of shown compounds as an energy source, electron acceptor, carbon source, or other nutrient source is indicated with different shapes. Note that compound concentrations, external supply, temperature, and pH are strongly site dependent. In continental serpentinizing systems, travertines form instead of chimneys. **(B)** Redox potential of available electron acceptors along a gradient from the oxic surface to the anoxic interior of the chimney wall. Adapted from Boyd et al. (2014).

Terrestrial and marine serpentinizing systems are likely to differ in environmental variables such as salinity and the compounds available for microbial metabolism, which might influence the ecosystem's overall functioning. An exciting transition site between terrestrial and marine serpentinite-hosted ecosystems is the Prony Bay Hydrothermal field located on the Southern Coast of New Caledonia, South Pacific (Launay and Fontes, 1985; Monnin et al., 2014). Prony Bay features several venting sites along a gradient from land to sea with a maximum depth of 50 m. The Prony Bay springs are fed by meteoric water, implying a strong salinity gradient between the hydrothermal fluid and ambient seawater (Monnin et al., 2014; Postec et al., 2015). Inversely, the Ney Springs system in Northern California features marine-type hydrothermal fluids in a continental context (Trutschel et al., 2022). Prony Bay is geochemically and microbiologically reflective of both ophiolitic and deep marine sites (Quéméneur et al., 2014; Postec et al., 2015; Frouin et al., 2018, 2022; Trutschel et al., 2022).

### 3. Metabolic strategies of serpentinite-hosted ecosystems

Serpentinite-hosted ecosystems are subjected to very challenging environmental conditions. The elevated pH poses a fundamental energetic problem on the cellular level, as it inverts the transmembrane pH gradient that typically drives all cellular processes. Furthermore, the high pH also reduces the bioavailability of electron acceptors, carbon, and other macronutrients (McCollom and Seewald, 2013; Schrenk et al., 2013). The microbial adaptations to these challenges are essential to understand life in a serpentinization context.

#### 3.1. The challenge of maintaining bioenergetics

##### 3.1.1. Maintaining pH homeostasis and a proton motive force

The central challenge for life in hyperalkaline conditions revolves around maintaining intracellular pH homeostasis and, more critically, conserving an electrochemical proton gradient across the cell membrane, which is the main driving force of the cell. The so-called proton motive force has two components: A transmembrane pH gradient ( $\Delta\text{pH}$ ), which is usually alkaline inside the cell relative to the outside, and a transmembrane electrical potential ( $\Delta\psi$ ), which is negative as long as the inner membrane surface is negatively charged. Maintaining a circum-neutral intracellular pH is crucial to ensure the stability of nucleic acids and proteins. In hyperalkaline conditions, however, the  $\Delta\text{pH}$  is inverted and very low due to  $\text{H}^+$  limitation outside of the cell, as the concentration of  $\text{H}^+$  decreases by  $10^{-4}$  at pH 11 compared to pH 7. This reduces the proton motive force and jeopardizes pH homeostasis (Krulwich et al., 2011 and references therein). Most alkaliphiles employ mechanisms increasing the efficiency of  $\text{H}^+$  uptake while maintaining a high  $\Delta\psi$ , which is also essential for pH homeostasis. This is achieved via  $\text{K}^+/\text{H}^+$  and  $\text{Na}^+/\text{H}^+$  antiporters with high  $\text{H}^+$  affinity, such as the Mrp complex in alkaliphilic *Bacillus* spp. (Ito et al., 2017). Those antiporters contribute to creating a transmembrane  $\text{Na}^+$  gradient, generating a sodium motive force that requires specialized sodium- $\text{F}_1\text{F}_0$ -ATP synthases or

$\text{Na}^+$ -dependent respiratory complexes (Krulwich et al., 2011; Kuhns et al., 2020). Based on varying  $\text{Na}^+$  concentrations, it is likely that these mechanisms differ between terrestrial and marine serpentinization-influenced site microorganisms. In low salt environments, alkaliphiles might excrete  $\text{Na}^+$  via  $\text{V}_1\text{V}_0$ -ATPases to maintain a sodium motive force (Suzuki et al., 2014; Ohlsson et al., 2019). Other alkaliphiles such as *Serpentinimonas* spp. isolated from The Cedars maintain a proton motive force using specialized  $\text{H}^+$  binding F type ATPases (Hicks et al., 2010; Suzuki et al., 2014).

##### 3.1.2. Energy conservation

The membrane potential generated via  $\text{H}^+$  or  $\text{Na}^+$  translocation depends on the redox potential of electron donors and acceptors. Serpentinization yields a range of reduced compounds that can serve as metabolic energy sources, most importantly  $\text{H}_2$  and  $\text{CH}_4$  (Figure 1A) (Boyd et al., 2014). The relative concentration of these gases varies significantly between sites (Etiope et al., 2011; Monnin et al., 2014), rendering generalized statements on a primary source of electrons provided by serpentinization difficult. While methanotrophs are more easily detected than hydrogenotrophs in many serpentinite-hosted environments (e.g., Brazelton et al., 2006; Kraus et al., 2021) (Table 1), the energetic potential of  $\text{H}_2$  oxidation greatly surpasses the oxidation potential of  $\text{CH}_4$ . The detection of hydrogenotrophs may be limited by the methodological approach, as the metabolic potential to oxidize  $\text{H}_2$  cannot be predicted from 16S rRNA sequences (Brazelton et al., 2012, 2022). Accordingly, metagenomic surveys show that many organisms in all types of serpentinite-hosted systems feature [FeFe]- and [NiFe]-hydrogenases (Brazelton et al., 2012; Mei et al., 2016; Kraus et al., 2021; Lecoeuvre et al., 2021; Frouin et al., 2022) (Table 1).

Another potential electron source in serpentinizing systems is constituted by compounds not directly created by serpentinization such as reduced sulfur species (Sabuda et al., 2020; Trutschel et al., 2022) and CO (Brazelton et al., 2012; Morrill et al., 2014; Fones et al., 2019) (Table 1). While the oxidation potential of CO is very low, the ability to use this energy source may provide a valuable ecological advantage. In surface exposed serpentinization-influenced waters, light constitutes an additional energy source used by cyanobacterial phototrophs (e.g., Kamran et al., 2020) (Table 1).

While serpentinization provides an abundance of electron donors, the availability of terminal electron acceptors is limited, especially in terrestrial serpentinizing systems, and mainly derived from the ambient environment (Figure 1A). Oxygen represents a very potent electron acceptor on the chimney or travertine surface. With increasing proximity to the reduced hydrothermal endmember, however, the availability of oxygen or alternative electron acceptors sharply decreases. The microbial community near the oxic-anoxic interphase may use nitrate (Frouin et al., 2022), even though data on nitrate reduction is scarce. In addition, organisms from the Troodos and The Cedars ophiolites have been shown to reduce metals such as iron or magnetite (Rizoulis et al., 2016; Rowe et al., 2017) (Table 1). Towards the anoxic interior of the hydrothermal carbonate chimneys or in deep ophiolite groundwaters, sulfate reduction is a dominant metabolic strategy in all types of serpentinite-hosted systems (Brazelton et al., 2006; Tiago and Verissimo, 2013; Postec et al., 2015; Glombitza et al., 2021) (Table 1 and Figure 1B).

To deal with electron acceptor limitation, many microbes also perform fermentation of sugars, simple organic acids and amino

**TABLE 1** Taxonomic survey for critical metabolic functions in all types of serpentinite-hosted environments, specifying the methodological approach including metagenome-assembled genomes (MAGs) and single-cell amplified genomes (SAGs).

Metabolism	Taxon	Site	Method	References
Hydrogen oxidation	Bacteria_Pseudomonadota_Alphaproteobacteria <sup>1</sup>	The Cedars (shallow waters)	MAGs	<a href="#">Suzuki et al. (2017)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria	The Cedars (shallow waters)	MAGs; 16S rRNA	<a href="#">Suzuki et al. (2013, 2017)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Betaproteobacteriales (formerly Betaproteobacteria) <sup>2</sup>	The Cedars (shallow waters)	MAGs	<a href="#">Suzuki et al. (2017)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Hydrogenophaga <sup>3</sup>	The Tablelands, Lost City (oxic/anoxic interface); Voltri; The Cedars (shallow waters); Prony Bay; Zambales	MAGs; 16S rRNA	<a href="#">Brazelton et al. (2012, 2017)</a> , <a href="#">Suzuki et al. (2013)</a> , <a href="#">Frouin et al. (2018)</a> , <a href="#">Woycheese et al. (2015)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas <sup>4</sup>	Voltri; The Cedars (shallow waters/non-specified)	16S rRNA; cultivation	<a href="#">Quéméneur et al. (2015)</a> , <a href="#">Suzuki et al. (2014, 2017)</a> , <a href="#">Brazelton et al. (2017)</a> , <a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. raichei <sup>5</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. barnesii <sup>6</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. maccroryi <sup>7</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Cupriavidus_C. necator (formerly <i>Ralstonia eutropha</i> ) <sup>8</sup>	The Tablelands, Lost City (oxic/anoxic interface)	MAGs	<a href="#">Brazelton et al. (2012)</a>
	Bacteria_Former class Deltaproteobacteria	The Cedars (shallow waters)	MAGs; 16S rRNA	<a href="#">Suzuki et al. (2013, 2017)</a>
	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales <sup>9</sup>	Voltri (deep subsurface)	MAGs	<a href="#">Brazelton et al. (2017)</a>
	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales_Desulfonatraceae_Desulfonatronum <sup>10</sup>	Prony Bay	16S rRNA	<a href="#">Postec et al. (2015)</a> , <a href="#">Mei et al. (2016)</a>
	Bacteria_BacillotaB_Desulfotomaculia_Desulfotomaculales_Desulfotomaculaceae_Desulfotomaculum_D. alkaliphilum <sup>11</sup>	Lost City	16S rRNA	<a href="#">Brazelton et al. (2006, 2010)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium_Lineage type I	Voltri; Samail (surface waters); Prony Bay	MAGs, SAGs, 14C labelling	<a href="#">Quéméneur et al. (2015, 2021, 2023)</a> , <a href="#">Fones et al. (2021)</a>
	Aerobic methane oxidation	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium_M. alkalithermotolerans strain DSM102889	La Crouen	Cultivation
Bacteria_Pseudomonadota_Gammaproteobacteria_Methylococcales_Methylococcaceae <sup>12</sup>		Voltri (shallow subsurface mixing zone)	16S rRNA, MAGs, 13C labelling	<a href="#">Brazelton et al. (2017)</a>
Bacteria_Pseudomonadota_Gammaproteobacteria_Methylococcales_Methylococcaceae_Methylococcus		Samail	16S rRNA	<a href="#">Kraus et al. (2021)</a>
Bacteria_Pseudomonadota_Alphaproteobacteria_Rhizobiales_Beijerinckiaceae_Methylosinus		Voltri	16S rRNA	<a href="#">Quéméneur et al. (2015)</a>

(Continued)



TABLE 1 (Continued)

Metabolism	Taxon	Site	Method	References
Methanogenesis/ anaerobic methane oxidation	Archaea_Halobacteriota_Methanomicrobia_Methanomicrobiales	Santa Elena	16S rRNA (MAGs)	<a href="#">Crespo-Medina et al. (2017)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales	Santa Elena; Voltri	16S rRNA (MAGs)	<a href="#">Crespo-Medina et al. (2017)</a> , <a href="#">Quéméneur et al. (2015)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae <sup>13</sup>	Voltri (deep subsurface)	16S rRNA, MAGs, 13C labelling	<a href="#">Brazelton et al. (2017)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium	Samail (subsurface); Voltri; Zambales	16S rRNA; MAGs; SAGs; 13C labelling; 14C labelling	<a href="#">Kraus et al. (2021)</a> , <a href="#">Brazelton et al. (2017)</a> , <a href="#">Fones et al. (2021)</a> , <a href="#">Woycheese et al. (2015)</a> , <a href="#">Quéméneur et al. (2015)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium_M. alcaliphilum strain DSM3387	Del Puerto	16S rRNA	<a href="#">Blank et al. (2009)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium_M. alkalithermotolerans strain DSM102889	La Crouen	Cultivation	<a href="#">Mei et al. (2022)</a>
	Archaea_Halobacteriota_Methanosarcinia_Methanosarcinales	Santa Elena; Voltri	16S rRNA (MAGs)	<a href="#">Crespo-Medina et al. (2017)</a> , <a href="#">Quéméneur et al. (2015)</a> , <a href="#">Suzuki et al. (2013)</a>
	Archaea_Halobacteriota_Methanosarcinia_Methanosarcinales_LCMS phylotype <sup>14</sup>	Lost City ( <a href="#">Brazelton et al. (2006)</a> ): High-temperature); Prony Bay (intertidal and submarine); Old City	16S rRNA, MAGs; 13C labelling	<a href="#">Schrenk et al. (2004)</a> , <a href="#">Brazelton et al. (2006, 2011)</a> , <a href="#">Frouin et al. (2018)</a> , <a href="#">Lecoeuvre et al. (2021)</a>
	Archaea_Halobacteriota_Methanosarcinia_Methanosarcinales_TCMS phylotype	The Cedars; Prony Bay (intertidal and submarine); Old City	16S rRNA, MAGs	<a href="#">Suzuki et al. (2013)</a> , <a href="#">Frouin et al. (2018)</a> , <a href="#">Lecoeuvre et al. (2021)</a>
	Archaea_Halobacteriota_Syntropharchaea_ANME-1	Lost City (low temperature); Santa Elena; Cabeço de Vide	16S rRNA (MAGs)	<a href="#">Brazelton et al. (2006)</a> , <a href="#">Crespo-Medina et al. (2017)</a> , <a href="#">Tiago and Verissimo (2013)</a>
Formate consumption	Bacteria_Pseudomonadota_Gammaproteobacteria_Methylococcales_Methylococcaceae <sup>12</sup>	Voltri (shallow subsurface mixing zone)	MAGs	<a href="#">Brazelton et al. (2017)</a>
	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales <sup>9</sup>	Voltri (deep subsurface)	MAGs	<a href="#">Brazelton et al. (2017)</a>
	Bacteria_Ca. Lithacetigena <sup>15</sup>	The Cedars, Hakuda Happo hot springs	MAGs	<a href="#">Nobu et al. (2022)</a>
	Bacteria_Ca. Bipolaricaulota (OP1/MSBL6) <sup>17</sup>	Lost City	MAGs	<a href="#">Brazelton et al. (2022)</a>
	Archaea_Methanobacteriota_Methanobacteria_Methanobacteriales_Methanobacteriaceae_Methanobacterium_Lineage type II	Samail (subsurface)	MAGs, SAGs, 14C labelling	<a href="#">Fones et al. (2021)</a>
Acetate consumption	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales <sup>9</sup>	Voltri (deep subsurface)	MAGs	<a href="#">Brazelton et al. (2017)</a>
	Archaea_Halobacteriota_Methanosarcinia_Methanosarcinales_LCMS phylotype <sup>14</sup>	Lost City	MAGs	<a href="#">Brazelton et al. (2011)</a>

(Continued)

TABLE 1 (Continued)

Metabolism	Taxon	Site	Method	References
Glycine consumption	Bacteria_Ca. Lithacetigena <sup>15</sup>	The Cedars, Hakuda Happo hot springs	MAGs	<a href="#">Nobu et al. (2022)</a>
Calcium carbonate consumption	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas <sup>4</sup>	The Cedars	Cultivation	<a href="#">Suzuki et al. (2014)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. raichei <sup>5</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. barnesii <sup>6</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Serpentinimonas_S. maccroryi <sup>7</sup>	The Cedars	Cultivation	<a href="#">Bird et al. (2021)</a>
	Bacteria_NPL-UPA2 clade <sup>18</sup>	The Cedars, Prony Bay, Lost City	MAGs	<a href="#">Suzuki et al. (2018)</a> , <a href="#">Brazelton et al. (2022)</a>
	Bacteria_Ca. Bipolaricaulota (OP1/MSBL6) <sup>17</sup>	Lost City	MAGs	<a href="#">Brazelton et al. (2022)</a>
	CO oxidation	Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Hydrogenophaga <sup>3</sup>	The Tablelands, Lost City (oxic/anoxic interface)	MAGs; 16S rRNA, 13C labelling
Bacteria_Pseudomonadota_Gammaproteobacteria_Burkholderiales_Burkholderiaceae_Cupriavidus_C. necator (formerly <i>Ralstonia eutropha</i> ) <sup>8</sup>		The Tablelands, Lost City (oxic/anoxic interface)	MAGs	<a href="#">Brazelton et al. (2012)</a>
Bacteria_Actinobacteriota_Ca. Hakubanella thermoalkaliphilus		Hakuda Happo hot springs	16S rRNA, SAGs	<a href="#">Merino et al. (2020)</a>
Archaea_Hadesarchaea (formerly SAGMEG)		Prony Bay	16S rRNA	<a href="#">Postec et al. (2015)</a>
Sulfide/sulfur oxidation	Bacteria_Pseudomonadota_Gammaproteobacteria_Betaproteobacteriales (formerly Betaproteobacteria) <sup>2</sup>	Cabeço de Vide	16S rRNA (DGGE)	<a href="#">Tiago and Verissimo (2013)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Thiomicrospirales_Thiomicrospiraceae_Hydrogenovibrio (formerly Thiomicrospira)	Lost City (low temperature)	16S rRNA	<a href="#">Brazelton et al. (2006)</a>
	Bacteria_Pseudomonadota_Alphaproteobacteria_Rhodobacterales_Rhodobacteraceae	Ney Springs	MAGs, Cultivation	<a href="#">Trutschel et al. (2022)</a>
	Bacteria_Pseudomonadota_Gammaproteobacteria_Pseudomonadales_Halomonadaceae_Halomonas	Ney Springs	MAGs, Cultivation	<a href="#">Trutschel et al. (2022)</a>

(Continued)

TABLE 1 (Continued)

Metabolism	Taxon	Site	Method	References
Sulfate reduction	Bacteria_BacillotaA_Clostridia <sup>16</sup>	Cabeço de Vide; The Cedars (deep subsurface)	16S rRNA (DGGE)	<a href="#">Tiago and Veríssimo (2013)</a> , <a href="#">Suzuki et al. (2013)</a>
	Bacteria_BacillotaB_Desulfotomaculia_Desulfotomaculales_Desulfotomaculaceae_Desulfotomaculum_D. alkaliphilum <sup>11</sup>	Lost City	16S rRNA	<a href="#">Brazelton et al. (2006, 2010)</a>
	Bacteria_BacillotaD_Dethiobacteria_Dethiobacterales_Dethiobacteraceae_Dethiobacter	The Tablelands, The Cedars, Cabeço de Vide, Prony Bay, Zambales		<a href="#">Brazelton et al. (2012)</a> , <a href="#">Suzuki et al. (2013)</a> , <a href="#">Tiago and Veríssimo (2013)</a> , <a href="#">Postec et al. (2015)</a> , <a href="#">Woycheese et al. (2015)</a> , <a href="#">Crespo-Medina et al. (2017)</a> , <a href="#">Trutschel et al. (2022)</a> , <a href="#">Twing et al. (2017)</a>
	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales <sup>9</sup>	Voltri	MAGs	<a href="#">Brazelton et al. (2017)</a>
	Bacteria_Desulfobacterota_Desulfovibrionia_Desulfovibrionales_Desulfonatronaceae_Desulfonatronum <sup>10</sup>	Prony Bay	16S rRNA	<a href="#">Postec et al. (2015)</a> , <a href="#">Mei et al. (2016)</a>
	Bacteria_Nitrospirota_Thermodesulfovibrionia_Thermodesulfovibrionales_Thermodesulfovibrionaceae	Samail (subsurface fluids)	16S rRNA	<a href="#">Rempfert et al. (2017)</a>
	Nitrogen fixation	Bacteria_Pseudomonadota_Alphaproteobacteria_Azospirillales_Azospirillaceae_Azospirillum	Voltri	16S rRNA
Archaea_Methanobacteriota_Methanobacteria_Methanobacterales_Methanobacteriaceae <sup>13</sup>		Voltri (deep subsurface)	16S rRNA, MAGs	<a href="#">Brazelton et al. (2017)</a>
Archaea_Halobacteriota_Methanosarcinia_Methanosarcinales_LCMS phylotype <sup>14</sup>		Lost City	MAGs	<a href="#">Brazelton et al. (2011)</a>
Phosphonate catabolism	Bacteria_Proteobacteria_Alphaproteobacteria <sup>1</sup>	Prony Bay, Lost City	MAGs	<a href="#">Frouin et al. (2022)</a>
	Bacteria_Proteobacteria_Gammaproteobacteria_Betaproteobacteriales (formerly Betaproteobacteria) <sup>2</sup>	Coast Range, Voltri, Santa Elena, Cabeço de Vide	MAGs	<a href="#">Frouin et al. (2022)</a>
	Bacteria_BacillotaA_Clostridia <sup>16</sup>	Prony Bay, Lost City	MAGs	<a href="#">Frouin et al. (2022)</a>
Acetogenesis	Bacteria_Chloroflexota (formerly Chloroflexi)	The Cedars (deep subsurface); Prony Bay (submarine)	MAGs; 16S rRNA	<a href="#">Suzuki et al. (2013, 2017)</a> , <a href="#">Frouin et al. (2018)</a>
	Bacteria_NPL-UPA2 clade <sup>18</sup>	The Cedars; Prony Bay	16S rRNA; MAGs	<a href="#">Postec et al. (2015)</a> , <a href="#">Mei et al. (2016)</a> , <a href="#">Suzuki et al. (2018)</a>
	Bacteria_BacillotaD_Dethiobacteria_Dethiobacterales_Dethiobacteraceae_Dethiobacter_D. alkaliphilus	The Cedars; Prony Bay; Cabeço de Vide	16S rRNA	<a href="#">Postec et al. (2015)</a> , <a href="#">Mei et al. (2016)</a> , <a href="#">Suzuki et al. (2013)</a> , <a href="#">Tiago and Veríssimo (2013)</a>
	Bacteria_Ca. Bipolaricaulota (OP1/MSBL6) <sup>17</sup>	Samail (subsurface fluids)	MAGs	<a href="#">Colman et al. (2022)</a>

(Continued)



TABLE 1 (Continued)

Metabolism	Taxon	Site	Method	References
Fermentation	Bacteria_BacillotaA_Clostridia	The Tablelands, Lost City	MAGs	<a href="#">Brazelton et al. (2012)</a>
	Bacteria_BacillotaA_Clostridia_Thermoanaerobacterales_Candidate Division OD1	The Cedars (deep subsurface); Voltri	16S rRNA, MAGs, 13C labelling	<a href="#">Suzuki et al. (2013, 2017)</a> , <a href="#">Brazelton et al. (2017)</a>
	Bacteria_BacillotaA_Clostridia_Lachnospirales_Vallitaleaceae_Vallitalea_V. pronyensis	Prony Bay	Cultivation	<a href="#">Ben Aissa et al. (2014)</a> , <a href="#">Mei et al. (2014)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Natronincolaceae_Alkaliphilus_A. hydrothermalis	Prony Bay	Cultivation	<a href="#">Ben Aissa et al. (2015)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Natronincolaceae_Alkaliphilus_A. serpentinus	Prony Bay	Cultivation	<a href="#">Postec et al. (2021)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Natronincolaceae_Alkaliphilus_A. pronyensis	Prony Bay	Cultivation	<a href="#">Postec et al. (2021)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Natronincolaceae_Serpentinicella_S. alkaliphila	Prony Bay	Cultivation	<a href="#">Mei et al. (2016)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Peptostreptococcaceae_Acetoanaerobium_A. pronyense	Prony Bay	Cultivation	<a href="#">Bes et al. (2015)</a>
	Bacteria_Bacteroidetes_Bacteroidia_Bacteroidales_ML635]-40	Voltri	16S rRNA, MAGs, 13C labelling	<a href="#">Brazelton et al. (2017)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Peptostreptococcaceae	Ney Springs	MAGs, 16S rRNA	<a href="#">Trutschel et al. (2022)</a>
	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Tindalliaceae_Tindallia	Ney Springs	MAGs, 16S rRNA	<a href="#">Trutschel et al. (2022)</a>
	Bacteria_BacillotaB_Desulfotomaculia_Desulfotomaculales_Desulfotomaculaceae_Desulfotomaculum	The Tablelands, Lost City	MAGs	<a href="#">Brazelton et al. (2012)</a>
	Bacteria_Bacilli_Bacillota_Erysipelotrichales_Erysipelotrichaceae	The Tablelands	MAGs	<a href="#">Brazelton et al. (2012)</a>
	Iron reduction	Bacteria_BacillotaA_Clostridia_Peptostreptococcales_Natronincolaceae_Alkaliphilus	Troodos	Cultivation
Bacteria_Bacillota_Bacilli_Paenibacillales_Paenibacillaceae_Paenibacillus		The Cedars	Cultivation	<a href="#">Rowe et al. (2017)</a>
Photosynthesis	Bacteria_Cyanobacteria_Cyanobacteriia_Leptolyngbyales_Leptolyngbyaceae_Leptolyngbya	Voltri (surface); Del Puerto	16S rRNA	<a href="#">Kamran et al. (2020)</a> , <a href="#">Blank et al. (2009)</a>
	Bacteria_Cyanobacteria_Cyanobacteriia_Synechococcales_Synechococcaceae_Synechococcus	Prony Bay	16S rRNA	<a href="#">Mei et al. (2016)</a>

Taxa appearing in several metabolic groups on the same taxonomic level are assigned with superscript numbers. The taxonomy is based on the Genome Taxonomy Database (GTDB Release 214).

acids, including Stickland type reactions (Barker, 1981; Postec et al., 2021). Therefore, a large proportion of the anaerobic serpentinite-hosted community may not feature a *bona fide* electron transport chain with cytochromes or quinones (Table 1). Genomic analysis suggests that many serpentinite-hosted fermenters conserve energy by substrate-level phosphorylation (e.g., in glycolysis), or via bifurcative-confurcative [FeFe] H<sub>2</sub>-producing hydrogenases, which balance the reducing equivalents NADH and ferredoxin produced by fermentation (Westphal et al., 2018). This is often associated with the Rnf complex, a respiratory enzyme that catalyzes the oxidation of reduced ferredoxin to the reduction of NAD<sup>+</sup>. The negative free energy change of this reaction is used to generate a transmembrane H<sup>+</sup> or Na<sup>+</sup> gradient (Westphal et al., 2018). This system can be considered a primitive respiratory mechanism where the terminal electron acceptor is H<sup>+</sup> (Buckel and Thauer, 2018).

## 3.2. The challenge of coping with nutrient limitation

Next to maintaining energy-yielding reactions, microorganisms in serpentinite-hosted environments must cope with severe nutrient limitation resulting from the decreased solubility of essential macronutrients at high pH (McCollom and Seewald, 2013; Schrenk et al., 2013). Especially relevant for the metabolic functioning of the community are the carbon, nitrogen, and phosphorous sources for primary production (Figure 1A).

### 3.2.1. Carbon sources and carbon fixation

One of the most significant issues regarding primary production in serpentinite-hosted environments is the absence of DIC, which precipitates as calcium carbonate in hyperalkaline conditions. While calcium carbonate is mostly insoluble and thus unavailable as a carbon source, it has nevertheless been shown to support the growth of some serpentinite-hosted microorganisms. This might be the result of local redissolution into bicarbonate catalyzed by the carbonic anhydrase (Suzuki et al., 2014; Fones et al., 2019; Bird et al., 2021) (Table 1). Alternatively, small organic molecules may serve as primary source of carbon. These include organic acids such as formate and acetate produced in Fischer-Tropsch and Sabatier-type reactions (Barbier et al., 2020; Fones et al., 2021) or via acetogenesis and fermentation (Kohl et al., 2016; Suzuki et al., 2017), as well as amino acids such as glycine produced in Strecker synthesis (Ménez et al., 2018; Nobu et al., 2022) (Figure 1A). The abiotic origin of those organic carbon sources tackles the definition of heterotrophy, which normally refers to the consumption of organic compounds derived from organic sources (Schönheit et al., 2016).

While bicarbonate, formate, acetate, and glycine have been shown to support the growth of microorganisms associated with serpentinitization, their metabolic route remains hypothetical. Transferred across the cell membrane via specialized transporters, formate can be oxidized to CO<sub>2</sub> via the formate dehydrogenase in the pH-neutral cytoplasm (Brazelton et al., 2022). Likewise, bicarbonate can be reduced to CO<sub>2</sub> via the carbonic anhydrase (Suzuki et al., 2014; Bird et al., 2021). The produced CO<sub>2</sub> is subsequently introduced to different carbon fixation pathways yielding acetyl-CoA. In serpentinite-hosted environments, the Wood-Ljungdahl pathway,

reverse tricarboxylic acid cycle, and Calvin-Benson-Bassham cycle have been confirmed (Seyler et al., 2020). Based on a recent study expanding the phylogenetic range of most carbon fixation pathways, the 3-hydroxypropionate bi-cycle, dicarboxylate/4-hydroxybutyrate cycle, and 3-hydroxypropionate/4-hydroxybutyrate cycle might also be employed (Garritano et al., 2022). Contrary to formate and bicarbonate, glycine can be directly transformed into acetyl-phosphate and subsequently acetyl-coA via the lesser known reductive glycine pathway (Sánchez-Andrea et al., 2020). Genes encoding the glycine reductase are found in metagenomes from Lost City, The Cedars, and the Japanese Hakuba Happo hot springs (Brazelton et al., 2022; Nobu et al., 2022). Also acetate can be directly transformed into acetyl-phosphate and acetyl-CoA, rendering its metabolic route less complex (Rose et al., 1954).

### 3.2.2. Sources of other nutrients

Serpentinization also decreases the solubility of other macronutrients essential for microbial growth. Inorganic phosphorous is severely limited in serpentinitizing environments because it is scavenged by the mineral brucite (Schrenk et al., 2013). A metagenomic survey revealed the high occurrence of genes involved in phosphonate catabolism in serpentinitizing sites, suggesting that the microbial community might use phosphonates as an alternative phosphorous source (Frouin et al., 2022). The catabolism of methylphosphonate, the most commonly available phosphonate species in marine environments, may additionally contribute to the global carbon and energy budget in these ecosystems by releasing CH<sub>4</sub> (Frouin et al., 2022). On the contrary, the availability of nitrogen in serpentinitizing environments remains controversial. While some authors suggest that concentrations are low (Schrenk et al., 2013), others propose that N<sub>2</sub> and nitrate are readily available to the serpentinite-hosted community (Lang et al., 2013; Rempfert et al., 2023). Potential nitrogen limitation may be alleviated by the fixation of N<sub>2</sub> derived from the endmember fluids or ambient seawater (Morrill et al., 2013; Monnin et al., 2014) (Figure 1A). While a recent study found the associated genetic marker *nifH* in 10 different serpentinite-hosted systems, its overall abundance was low (Frouin et al., 2022).

## 3.3. Metabolic links to the emergence of life

The biochemical characteristics of serpentinite-hosted ecosystems reinforce the presumed link between serpentinitization and the beginnings of life. Serpentinization is an ancient process which likely occurred on early Earth (Russell et al., 2010). The abiotic production of organic acids associated with serpentinitization is for instance supported by isotopic signatures (McCollom and Seewald, 2013). Likewise, amino acids such as glycine may be formed abiotically, which is especially interesting in prebiotic chemistry (Aubrey et al., 2009; Ménez et al., 2018). The metabolic use of these compounds is linked to very deep-branching functions, such as the reductive tricarboxylic acid cycle and the Wood-Ljungdahl pathway, which are likely the most ancient carbon fixation pathways on Earth (Sumi and Harada, 2021). Another primordial function preserved in serpentinitizing environments is CO oxidation. CO is not only one of the most ancient energy sources exploited in metabolism but is also

suggested to have played a key role in several critical prebiotic reactions (King and Weber, 2007). It may thus constitute a direct link between abiotic and biotic chemistry.

## 4. Diversity of identified metabolic key players

The specific metabolic challenges posed by serpentinization suggest the presence of specialized taxonomic groups playing an important role in the trophic network. It might be possible that such “core” taxa are relevant in a wide variety of serpentinizing environments, even though the overall community structure can vary significantly in space and time (Suzuki et al., 2013; Postec et al., 2015; Fones et al., 2019; Brazelton et al., 2022; Trutschel et al., 2022).

The community of H<sub>2</sub> oxidizers appears to be dominated by Gammaproteobacteria (Table 1). This includes a major proportion of *Serpentinimonas* (formerly grouped under Betaproteobacteria). *Serpentinimonas* is one of the taxa most commonly associated with serpentinization, and represented by some of the few available isolates from serpentinite-hosted ecosystems (Suzuki et al., 2014; Bird et al., 2021). So far, all of those isolated strains originate from The Cedars. Still, 16S rRNA analysis confirms the presence of *Serpentinimonas* and its sister genus *Hydrogenophaga* in other terrestrial systems, as well as in Prony Bay and Lost City (e.g., Brazelton et al., 2012; Quéménéur et al., 2015; Woycheese et al., 2015; Frouin et al., 2018) (Table 1).

While aerobic methane oxidation is mostly performed by the bacterial *Methylococcales* (Brazelton et al., 2017; Kraus et al., 2021), anaerobic methanotrophy and methanogenesis feature exclusively archaea (Table 1). Hydrogenotrophic methanogens belonging to *Methanobacteriales* are often detected in serpentinite-hosted terrestrial ecosystems (Woycheese et al., 2015; Brazelton et al., 2017; Kraus et al., 2021; Quéménéur et al., 2021, 2023; Mei et al., 2022). In addition, there is a subgroup of *Methanosarcinales* which is probably endemic to serpentinizing environments and includes two distinct phylotypes (e.g., Schrenk et al., 2004; Brazelton et al., 2010; Suzuki et al., 2013). Next to their systems of origin, The Cedars *Methanosarcinales* (TCMS) and Lost City *Methanosarcinales* (LCMS) have been observed in Prony Bay (Frouin et al., 2018) and Old City (Lecoeuvre et al., 2021) (Table 1). However, attempts to culture them have been unsuccessful so far.

The fermenting community seems almost entirely dominated by *Clostridia* (Table 1), of which several novel species have been isolated from Prony Bay (Ben Aissa et al., 2014, 2015; Mei et al., 2014; Bes et al., 2015; Postec et al., 2021).

Other energy yielding metabolic strategies including CO oxidation, sulfur oxidation and sulfate reduction are performed by a broader diversity of taxonomic groups (Table 1). Notably, the community of sulfate reducers includes *Desulfovibrionales* and *Dethiobacter* species, which can be very abundant in serpentinizing environments (Brazelton et al., 2012; Suzuki et al., 2013; Tiago and Verissimo, 2013; Postec et al., 2015; Woycheese et al., 2015; Mei et al., 2016) (Table 1).

Regarding carbon uptake, it might be especially interesting to further investigate certain candidate phyla that occur in various serpentinizing systems. For example, *Ca. Bipolaricaulota* can use bicarbonate and formate (Brazelton et al., 2022) and plays a role in acetogenesis (Colman et al., 2022). Likewise, *Ca. NPL-UPA2* grows

on bicarbonate (Brazelton et al., 2022) and performs acetogenesis via the Wood-Ljungdahl pathway (Suzuki et al., 2018) (Table 1). The Wood-Ljungdahl pathway is also employed by *Ca. Hakubanella thermoalkaliphilus*, a novel Actinobacteriota from the Hakuba Happo hot springs serpentinizing system (Merino et al., 2020). Finally, *Ca. Lithacetigena* was recently shown to perform glycine reduction (Nobu et al., 2022). These candidate phyla might play an important role in the trophic chain by supplying fixed carbon to the community.

## 5. Methodological shortcomings and future challenges

While several critical metabolic strategies and taxonomic groups could be identified, the scope of their distribution across serpentinizing ecosystems remains unclear due to methodological biases and shortcomings. Firstly, continental sites have been studied much more extensively than marine ones (Table 1). Our understanding of the latter is almost entirely based on the famous Lost City (Table 1) and most recently Old City (Lecoeuvre et al., 2021), which limits the generalization of findings on marine serpentinizing systems and reduces the meaningfulness of comparison with continental ones. This issue emphasizes the interest of the shallow marine transition field of Prony Bay. Its common characteristics with continental and marine sites may help establish the core metabolic properties of serpentinite-hosted ecosystems. In addition, the study of such shallow fields is facilitated by their geographical accessibility.

Another factor introducing bias is the methodological approach. Most studies rely on metabarcoding and metagenomic techniques (Table 1), which are strongly dependent on the scope of available reference databases. Moreover, the presence of a functional gene does not necessarily signify its activity. However, confirmation of gene expression is rare, notably due to technical difficulties in obtaining quality metatranscriptomes from such environments (Table 1). In addition, there is a lack of experimental evidence complementing bioinformatic hypotheses. Studies attempting to bridge this gap include activity measurements using <sup>13</sup>C and <sup>14</sup>C labeled substrates in microcosms (Brazelton et al., 2011, 2017; Morrill et al., 2014; Fones et al., 2021), as well as isolation of *Serpentinimonas* and *Clostridia* species from cultures (e.g., Suzuki et al., 2014; Postec et al., 2021) (Table 1). Their small number is probably also associated with technical difficulties, including the cultivation of recalcitrant microorganisms such as obligate anaerobic chemolithoautotrophs. While combined omics approaches can provide valuable results, critical metabolic groups will eventually need to be cultivated to confirm their functional role in the ecosystem. This may be facilitated by implementing more sophisticated culture platforms to mimic the conditions associated with serpentinization in the laboratory.

## 6. Conclusion

Serpentinite-hosted environments are inhabited by microbial communities that cope with energetic challenges and severe nutrient limitation. It can be assumed that a significant proportion of those microorganisms yield energy from H<sub>2</sub> oxidation with electron acceptors derived from external sources or CO<sub>2</sub> degassed from mantle rocks. Calcium carbonate can serve as inorganic carbon source, and

formate, acetate and glycine as organic carbon sources for primary production. Bicarbonate, formate, and glycine may be fixed via different carbon fixation pathways such as the Wood-Ljungdahl pathway, the reverse tricarboxylic acid and Calvin-Benson-Bassham cycles and the reductive glycine pathway. Moreover, the microbial communities might cope with nitrogen and phosphorous limitation by fixing  $N_2$  and breaking down phosphonates. The analysis of functional genes suggests that taxa such as *Gammaproteobacteria*, *Desulfovibrionales*, *Clostridia* and several candidate phyla play a crucial role in the trophic network and that the genera *Serpentinimonas*, *Hydrogenophaga*, and *Methanobacterium* as well as uncultivated *Methanosarcinales*, are characteristic for serpentinizing environments. However, the scope of research on serpentinite-hosted ecosystems needs to be broadened by including a greater diversity of marine and shallow transition sites. In addition, experimental evidence is needed to confirm the metabolic activity of hypothesized key players. A technological advancement of the methodological approach might not only contribute to the understanding of present serpentinite-hosted ecosystems, but also provide insights into the beginning of life on Earth and potentially elsewhere.

## Author contributions

GE, AP, and RP: conceptualization. AL and MQ: validation. RP: investigation and writing of original draft. AL, MQ, AP, GE, and RP: review and edit of original draft. GE and AP: supervision. GE: project administration. All authors contributed to the article and approved the submitted version.

## References

- Albers, E., Bach, W., Pérez-Gussinyé, M., McCammon, C., and Frederichs, T. (2021). Serpentinization-driven  $H_2$  production from continental break-up to mid-ocean ridge spreading: unexpected high rates at the West Iberia margin. *Front. Earth Sci.* 9:673063. doi: 10.3389/feart.2021.673063
- Aubrey, A. D., Cleaves, H. J., and Bada, J. L. (2009). The role of submarine hydrothermal systems in the synthesis of amino acids. *Orig. Life Evol. Biosph.* 39, 91–108. doi: 10.1007/s11084-008-9153-2
- Barbier, S., Huang, F., Andreani, M., Tao, R., Hao, J., Eleish, A., et al. (2020). A review of  $H_2$ ,  $CH_4$ , and hydrocarbon formation in experimental serpentinization using network analysis. *Front. Earth Sci.* 8:209. doi: 10.3389/feart.2020.00209
- Barker, H. A. (1981). Amino acid degradation by anaerobic bacteria. *Annu. Rev. Biochem.* 50, 23–40. doi: 10.1146/annurev.bi.50.070181.000323
- Barnes, I., O'Neil, J. R., and Trescases, J. J. (1978). Present day serpentinization in New Caledonia, Oman and Yugoslavia. *Geochim. Cosmochim. Acta* 42, 144–145. doi: 10.1016/0016-7037(78)90225-9
- Ben Aissa, F., Postec, A., Erauso, G., Payri, C., Pelletier, B., Hamdi, M., et al. (2015). Characterization of *Alkaliphilus hydrothermalis* sp. nov., a novel alkaliphilic anaerobic bacterium, isolated from a carbonaceous chimney of the Prony hydrothermal field, New Caledonia. *Extremophiles* 19, 183–188. doi: 10.1007/s00792-014-0697-y
- Ben Aissa, F., Postec, A., Erauso, G., Payri, C., Pelletier, B., Hamdi, M., et al. (2014). *Vallitalea pronyensis* sp. nov., isolated from a marine alkaline hydrothermal chimney. *Int. J. Syst. Evol. Microbiol.* 64, 1160–1165. doi: 10.1099/ijs.0.055756-0
- Bes, M., Merrouch, M., Joseph, M., Quéméneur, M., Payri, C., Pelletier, B., et al. (2015). *Acetoanaerobium pronyense* sp. nov., an anaerobic alkaliphilic bacterium isolated from a carbonate chimney of the Prony hydrothermal field (New Caledonia). *Int. J. Syst. Evol. Microbiol.* 65, 2574–2580. doi: 10.1099/ijs.0.000307
- Bird, L. J., Kuenen, J. G., Osburn, M. R., Tomioka, N., Ishii, S., Barr, C., et al. (2021). *Serpentinimonas* gen. Nov., *Serpentinimonas raichei* sp. nov., *Serpentinimonas barnesii* sp. nov. and *Serpentinimonas maccrorryi* sp. nov., hyperalkaliphilic and facultative autotrophic bacteria isolated from terrestrial serpentinizing springs. *Int. J. Syst. Evol. Microbiol.* 71:004945. doi: 10.1099/ijsem.0.004945
- Blank, J. G., Green, S. J., Blake, D., Valley, J. W., Kita, N. T., Treiman, A., et al. (2009). An alkaline spring system within the Del Puerto ophiolite (California, USA): a Mars analog site. *Planet. Space Sci.* 57, 533–540. doi: 10.1016/j.pss.2008.11.018
- Boyd, E. S., Schut, G. J., Peters, J. W., and Adams, M. W. W. (2014). Hydrogen metabolism and the evolution of biological respiration: two separate families of enzymes that oxidize hydrogen and also produce it arose through convergent evolution. *Microbe Mag.* 9, 361–367. doi: 10.1128/microbe.9.361.1
- Brazelton, W. J., Ludwig, K. A., Sogin, M. L., Andreishcheva, E. N., Kelley, D. S., Shen, C.-C., et al. (2010). Archaea and bacteria with surprising microdiversity show shifts in dominance over 1,000-year time scales in hydrothermal chimneys. *Proc. Natl. Acad. Sci. USA* 107, 1612–1617. doi: 10.1073/pnas.0905369107
- Brazelton, W. J., McGonigle, J. M., Motamedi, S., Pendleton, H. L., Twing, K. L., Miller, B. C., et al. (2022). Metabolic strategies shared by basement residents of the Lost City hydrothermal field. *Appl. Environ. Microbiol.* 88:e0092922. doi: 10.1128/aem.00929-22
- Brazelton, W. J., Mehta, M. P., Kelley, D. S., and Baross, J. A. (2011). Physiological differentiation within a single-species biofilm fueled by serpentinization. *mBio* 2, e00127–e00111. doi: 10.1128/mBio.00127-11
- Brazelton, W. J., Morrill, P. L., Szponar, N., and Schrenk, M. O. (2013). Bacterial communities associated with subsurface geochemical processes in continental serpentinite springs. *Appl. Environ. Microbiol.* 79, 3906–3916. doi: 10.1128/AEM.00330-13
- Brazelton, W. J., Nelson, B., and Schrenk, M. O. (2012). Metagenomic evidence for  $H_2$  oxidation and  $H_2$  production by serpentinite-hosted subsurface microbial communities. *Front. Microbiol.* 2:268. doi: 10.3389/fmicb.2011.00268
- Brazelton, W. J., Schrenk, M. O., Kelley, D. S., and Baross, J. A. (2006). Methane- and sulfur-metabolizing microbial communities dominate the Lost City hydrothermal field ecosystem. *Appl. Environ. Microbiol.* 72, 6257–6270. doi: 10.1128/AEM.00574-06
- Brazelton, W. J., Thornton, C. N., Hyer, A., Twing, K. L., Longino, A. A., Lang, S. Q., et al. (2017). Metagenomic identification of active methanogens and methanotrophs in serpentinite springs of the Voltri massif, Italy. *PeerJ* 5:e2945. doi: 10.7717/peerj.2945

## Funding

This project was financially supported by the ANR MICROPRONY (N°19-CE02-0020-02), the French Institute of Research for Development (IRD), and a Ph.D. fellowship granted to RP by the Aix-Marseille University's Doctoral School "Sciences de l'Environnement" (ED 251).

## Acknowledgments

The authors thank all project partners, including MICROPRONY.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



- Buckel, W., and Thauer, R. K. (2018). Flavin-based electron bifurcation, ferredoxin, flavodoxin, and anaerobic respiration with protons (Ech) or NAD<sup>+</sup> (Rnf) as electron acceptors: a historical review. *Front. Microbiol.* 9:401. doi: 10.3389/fmicb.2018.00401
- Colman, D. R., Kraus, E. A., Thieringer, P. H., Rempfert, K., Templeton, A. S., Spear, J. R., et al. (2022). Deep-branching acetogens in serpentinized subsurface fluids of Oman. *Proc. Natl. Acad. Sci. USA* 119:e2206845119. doi: 10.1073/pnas.2206845119
- Crespo-Medina, M., Twing, K. I., Sánchez-Murillo, R., Brazelton, W. J., McCollom, T. M., and Schrenk, M. O. (2017). Methane dynamics in a tropical serpentinizing environment: the Santa Elena ophiolite, Costa Rica. *Front. Microbiol.* 8:916. doi: 10.3389/fmicb.2017.00916
- Daas, F. L., Økland, I., Dahle, H., Jørgensen, S. L., Thorseth, I. H., and Pedersen, R. B. (2013). Microbial life associated with low-temperature alteration of ultramafic rocks in the Leka ophiolite complex. *Geobiology* 11, 318–339. doi: 10.1111/gbi.12035
- Etiopio, G., Schoell, M., and Hosgörmez, H. (2011). Abiotic methane flux from the Chimaera seep and Tekirova ophiolites (Turkey): understanding gas exhalation from low temperature serpentinization and implications for Mars. *Earth Planet. Sci. Lett.* 310, 96–104. doi: 10.1016/j.epsl.2011.08.001
- Fabri, M.-C., Bargain, A., Briand, P., Gebruk, A., Fouquet, Y., Morineaux, M., et al. (2011). The hydrothermal vent community of a new deep-sea field, Ashadze-1, 12°58'N on the mid-Atlantic ridge. *J. Mar. Biol. Assoc. U. K.* 91, 1–13. doi: 10.1017/S0025315410000731
- Flores, G. E., Campbell, J. H., Kirshtein, J. D., Meneghin, J., Podar, M., Steinberg, J. I., et al. (2011). Microbial community structure of hydrothermal deposits from geochemically different vent fields along the mid-Atlantic ridge: microbial communities of hydrothermal vent deposits. *Environ. Microbiol.* 13, 2158–2171. doi: 10.1111/j.1462-2920.2011.02463.x
- Fones, E. M., Colman, D. R., Kraus, E. A., Nothaft, D. B., Poudel, S., Rempfert, K. R., et al. (2019). Physiological adaptations to serpentinization in the Samail ophiolite, Oman. *ISME J.* 13, 1750–1762. doi: 10.1038/s41396-019-0391-2
- Fones, E. M., Colman, D. R., Kraus, E. A., Stepanauskas, R., Templeton, A. S., Spear, J. R., et al. (2021). Diversification of methanogens into hyperalkaline serpentinizing environments through adaptations to minimize oxidant limitation. *ISME J.* 15, 1121–1135. doi: 10.1038/s41396-020-00838-1
- Frouin, E., Bes, M., Ollivier, B., Quéméneur, M., Postec, A., Debros, D., et al. (2018). Diversity of rare and abundant prokaryotic phylotypes in the Prony hydrothermal field and comparison with other serpentinite-hosted ecosystems. *Front. Microbiol.* 9:102. doi: 10.3389/fmicb.2018.00102
- Frouin, E., Lecoivre, A., Armougom, F., Schrenk, M. O., and Erauso, G. (2022). Comparative metagenomics highlight a widespread pathway involved in catabolism of phosphonates in marine and terrestrial serpentinizing. *Ecosystems* 7:e0032822. doi: 10.1128/msystems.00328-22
- Früh-Green, G. L., Connolly, J. A. D., Plas, A., Kelley, D. S., and Grobety, B. (2004). “Serpentinization of oceanic peridotites: implications for geochemical cycles and biological activity” in *Geophysical monograph series*. eds. W. S. D. Wilcock, E. F. DeLong, D. S. Kelley, J. A. Baross and S. Craig Cary (Washington, D C: American Geophysical Union), 119–136. doi: 10.1029/144GM08
- Garritano, A. N., Song, W., and Thomas, T. (2022). Carbon fixation pathways across the bacterial and archaeal tree of life. *PNAS Nexus* 1:pgac226. doi: 10.1093/pnasnexus/pgac226
- Glombitza, C., Putman, L. I., Rempfert, K. R., Kubo, M. D., Schrenk, M. O., Templeton, A. S., et al. (2021). Active microbial sulfate reduction in fluids of serpentinizing peridotites of the continental subsurface. *Commun. Earth Environ.* 2:84. doi: 10.1038/s43247-021-00157-z
- Hicks, D. B., Liu, J., Fujisawa, M., and Krulwich, T. A. (2010). F1F0-ATP synthases of alkaliphilic bacteria: lessons from their adaptations. *Biochim. Biophys. Acta BBA – Bioenerg.* 1797, 1362–1377. doi: 10.1016/j.bbabo.2010.02.028
- Ito, M., Morino, M., and Krulwich, T. A. (2017). Mrp antiporters have important roles in diverse bacteria and archaea. *Front. Microbiol.* 8:2325. doi: 10.3389/fmicb.2017.02325
- Kamran, A., Sauter, K., Reimer, A., Wacker, T., Reitner, J., and Hoppert, M. (2020). Cyanobacterial mats in calcite-precipitating serpentinite-hosted alkaline springs of the Voltri Massif, Italy. *Microorganisms* 9:62. doi: 10.3390/microorganisms9010062
- Kelley, D. S., Karson, J. A., Früh-Green, G. L., Yoerger, D. R., Shank, T. M., Butterfield, D. A., et al. (2005). A serpentinite-hosted ecosystem: the Lost City hydrothermal field. *Science* 307, 1428–1434. doi: 10.1126/science.1102556
- King, G. M., and Weber, C. F. (2007). Distribution, diversity and ecology of aerobic CO-oxidizing bacteria. *Nat. Rev. Microbiol.* 5, 107–118. doi: 10.1038/nrmicro1595
- Kohl, L., Cumming, E., Cox, A., Rietze, A., Morrissey, L., Lang, S. Q., et al. (2016). Exploring the metabolic potential of microbial communities in ultra-basic, reducing springs at the cedars, CA, USA: experimental evidence of microbial methanogenesis and heterotrophic acetogenesis: methanogenesis at the cedars. *J. Geophys. Res. Biogeosci.* 121, 1203–1220. doi: 10.1002/2015JG003233
- Kraus, E. A., Nothaft, D., Stamps, B. W., Rempfert, K. R., Ellison, E. T., Matter, J. M., et al. (2021). Molecular evidence for an active microbial methane cycle in subsurface serpentinite-hosted groundwaters in the Samail ophiolite, Oman. *Appl. Environ. Microbiol.* 87, e02068–e02020. doi: 10.1128/AEM.02068-20
- Krulwich, T. A., Sachs, G., and Padan, E. (2011). Molecular aspects of bacterial pH sensing and homeostasis. *Nat. Rev. Microbiol.* 9, 330–343. doi: 10.1038/nrmicro2549
- Kuhns, M., Trifunović, D., Huber, H., and Müller, V. (2020). The Rnf complex is a Na<sup>+</sup> coupled respiratory enzyme in a fermenting bacterium, *Thermotoga maritima*. *Commun. Biol.* 3:431. doi: 10.1038/s42003-020-01158-y
- Lane, N. (2015). *The vital question: energy, evolution, and the origins of complex life*. New York City: WW Norton & Company.
- Lang, S. Q., Früh-Green, G. L., Bernasconi, S. M., and Butterfield, D. A. (2013). Sources of organic nitrogen at the serpentinite-hosted Lost City hydrothermal field. *Geobiology* 11, 154–169. doi: 10.1111/gbi.12026
- Launay, J., and Fontes, J.-C. (1985). Les sources thermales de Prony (Nouvelle Calédonie) et leurs précipités chimiques, exemple de formation de brucite primaire. *Geol. Fr.* 1, 83–100.
- Lecoivre, A., Ménez, B., Cannat, M., Chavagnac, V., and Gérard, E. (2021). Microbial ecology of the newly discovered serpentinite-hosted Old City hydrothermal field (southwest Indian ridge). *ISME J.* 15, 818–832. doi: 10.1038/s41396-020-00816-7
- McCollom, T. M., and Seewald, J. S. (2001). A reassessment of the potential for reduction of dissolved CO<sub>2</sub> to hydrocarbons during serpentinization of olivine. *Geochim. Cosmochim. Acta* 65, 3769–3778. doi: 10.1016/S0016-7037(01)00655-X
- McCollom, T. M., and Seewald, J. S. (2013). Serpentinites, hydrogen, and life. *Elements* 9, 129–134. doi: 10.2113/gselements.9.2.129
- Mei, N., Postec, A., Bartoli, M., Vandecasteele, C., Wils, L., Gil, L., et al. (2022). Methanobacterium alkalithermotolerans sp. nov., a novel alkaliphilic and hydrogen-utilizing methanogen isolated from an alkaline geothermal spring (La Crouen, New Caledonia). *Int. J. Syst. Evol. Microbiol.* 72. doi: 10.1099/ijsem.0.005554
- Mei, N., Postec, A., Monnin, C., Pelletier, B., Payri, C. E., Ménez, B., et al. (2016). Metagenomic and PCR-based diversity surveys of [FeFe]-hydrogenases combined with isolation of alkaliphilic hydrogen-producing bacteria from the serpentinite-hosted Prony hydrothermal field, New Caledonia. *Front. Microbiol.* 7:1301. doi: 10.3389/fmicb.2016.01301
- Mei, N., Zergane, N., Postec, A., Erauso, G., Ollier, A., Payri, C., et al. (2014). Fermentative hydrogen production by a new alkaliphilic Clostridium sp. (strain PROH2) isolated from a shallow submarine hydrothermal chimney in Prony Bay, New Caledonia. *Int. J. Hydrog. Energy* 39, 19465–19473. doi: 10.1016/j.ijhydene.2014.09.111
- Ménez, B., Pisapia, C., Andreani, M., Jamme, F., Vanbellingen, Q. P., Brunelle, A., et al. (2018). Abiotic synthesis of amino acids in the recesses of the oceanic lithosphere. *Nature* 564, 59–63. doi: 10.1038/s41586-018-0684-z
- Merino, N., Kawai, M., Boyd, E. S., Colman, D. R., McGlynn, S. E., Neelson, K. H., et al. (2020). Single-cell genomics of novel actinobacteria with the Wood-Ljungdahl pathway discovered in a serpentinizing system. *Front. Microbiol.* 11:1031. doi: 10.3389/fmicb.2020.01031
- Monnin, C., Chavagnac, V., Boulart, C., Ménez, B., Gérard, M., Gérard, E., et al. (2014). Fluid chemistry of the low temperature hyperalkaline hydrothermal system of Prony Bay (New Caledonia). *Biogeosciences* 11, 5687–5706. doi: 10.5194/bg-11-5687-2014
- Morrill, P. L., Brazelton, W. J., Kohl, L., Rietze, A., Miles, S. M., Kavanagh, H., et al. (2014). Investigations of potential microbial methanogenic and carbon monoxide utilization pathways in ultra-basic reducing springs associated with present-day continental serpentinization: the tablelands, NL, CAN. *Front. Microbiol.* 5:613. doi: 10.3389/fmicb.2014.00613
- Morrill, P. L., Kuonen, J. G., Johnson, O. J., Suzuki, S., Rietze, A., Sessions, A. L., et al. (2013). Geochemistry and geobiology of a present-day serpentinization site in California: the cedars. *Geochim. Cosmochim. Acta* 109, 222–240. doi: 10.1016/j.gca.2013.01.043
- Nakamura, K., Morishita, T., Bach, W., Klein, F., Hara, K., Okino, K., et al. (2009). Serpentinized troctolites exposed near the Kairei hydrothermal field, central Indian ridge: insights into the origin of the Kairei hydrothermal fluid supporting a unique microbial ecosystem. *Earth Planet. Sci. Lett.* 280, 128–136. doi: 10.1016/j.epsl.2009.01.024
- Neubeck, A., Sun, L., Müller, B., Ivarsson, M., Hosgörmez, H., Özcan, D., et al. (2017). Microbial community structure in a serpentine-hosted abiotic gas seepage at the Chimaera ophiolite, Turkey. *Appl. Environ. Microbiol.* 83, e03430–e03416. doi: 10.1128/AEM.03430-16
- Nobu, M. K., Nakai, R., Tamazawa, S., Mori, H., Toyoda, A., Ijiri, A., et al. (2022). Unique H<sub>2</sub>-utilizing lithotrophy in serpentinite-hosted systems. *ISME J.* 17, 95–104. doi: 10.1038/s41396-022-01197-9
- Ohlsson, J. I., Osvatic, J. T., Becraft, E. D., and Swingley, W. D. (2019). Microbial community in hyperalkaline steel slag-fill emulates serpentinizing springs. *Diversity* 11:103. doi: 10.3390/d11070103
- Perner, M., Kuever, J., Seifert, R., Pape, T., Koschinsky, A., Schmidt, K., et al. (2007). The influence of ultramafic rocks on microbial communities at the Logatchev hydrothermal field, located 15°N on the mid-Atlantic ridge: influence of ultramafic rocks on microbial communities. *FEMS Microbiol. Ecol.* 61, 97–109. doi: 10.1111/j.1574-6941.2007.00325.x
- Postec, A., Quéméneur, M., Bes, M., Mei, N., Benaïssa, F., Payri, C., et al. (2015). Microbial diversity in a submarine carbonate edifice from the serpentinizing

- hydrothermal system of the Prony Bay (New Caledonia) over a 6-year period. *Front. Microbiol.* 6:857. doi: 10.3389/fmicb.2015.00857
- Postec, A., Quéméneur, M., Lecoeuvre, A., Chabert, N., Joseph, M., and Erauso, G. (2021). Alkaliphilus serpentinus sp. nov. and Alkaliphilus pronyensis sp. nov., two novel anaerobic alkaliphilic species isolated from the serpentinite-hosted Prony Bay hydrothermal field (New Caledonia). *Syst. Appl. Microbiol.* 44:126175. doi: 10.1016/j.syapm.2020.126175
- Quéméneur, M., Bes, M., Postec, A., Mei, N., Hamelin, J., Monnin, C., et al. (2014). Spatial distribution of microbial communities in the shallow submarine alkaline hydrothermal field of the Prony Bay, New Caledonia: microbial communities of Prony hydrothermal field. *Environ. Microbiol. Rep.* 6, 665–674. doi: 10.1111/1758-2229.12184
- Quéméneur, M., Mei, N., Monnin, C., Postec, A., Guasco, S., Jeanpert, et al. (2023). Microbial taxa related to natural hydrogen and methane emissions in serpentinite-hosted hyperalkaline springs of New Caledonia. *Front. Microbiol.* 14:1196516. doi: 10.3389/fmicb.2023.1196516
- Quéméneur, M., Mei, N., Monnin, C., Postec, A., Wils, L., Bartoli, M., et al. (2021). Prokaryotic diversity and hydrogenotrophic methanogenesis in an alkaline spring (La Crouen, New Caledonia). *Microorganisms* 9:1360. doi: 10.3390/microorganisms9071360
- Quéméneur, M., Palvadeau, A., Postec, A., Monnin, C., Chavagnac, V., Ollivier, B., et al. (2015). Endolithic microbial communities in carbonate precipitates from serpentinite-hosted hyperalkaline springs of the Voltri massif (Ligurian Alps, northern Italy). *Environ. Sci. Pollut. Res.* 22, 13613–13624. doi: 10.1007/s11356-015-4113-7
- Rempfert, K. R., Miller, H. M., Bompard, N., Nothaft, D., Matter, J. M., Kelemen, P., et al. (2017). Geological and geochemical controls on subsurface microbial life in the Samail ophiolite, Oman. *Front. Microbiol.* 8:56. doi: 10.3389/fmicb.2017.00056
- Rempfert, K. R., Nothaft, D. B., Kraus, E. A., Asamoto, C. K., Evans, R. D., Spear, J. R., et al. (2023). Subsurface biogeochemical cycling of nitrogen in the actively serpentinizing Samail ophiolite, Oman. *Front. Microbiol.* 14:1139633. doi: 10.3389/fmicb.2023.1139633
- Rizoulis, A., Milodowski, A. E., Morris, K., and Lloyd, J. R. (2016). Bacterial diversity in the hyperalkaline allas springs (Cyprus), a natural analogue for cementitious radioactive waste repository. *Geomicrobiol. J.* 33, 73–84. doi: 10.1080/01490451.2014.961107
- Rose, I. A., Grunberg-Manago, M., Korey, S. R., and Ochoa, S. (1954). Enzymatic phosphorylation of acetate. *J. Biol. Chem.* 211, 737–756. doi: 10.1016/S0021-9258(18)71161-7
- Rowe, A. R., Yoshimura, M., LaRowe, D. E., Bird, L. J., Amend, J. P., Hashimoto, K., et al. (2017). In situ electrochemical enrichment and isolation of a magnetite-reducing bacterium from a high pH serpentinizing spring. *Environ. Microbiol.* 19, 2272–2285. doi: 10.1111/1462-2920.13723
- Russell, M. J., Hall, A. J., Cairns-Smith, A. G., and Braterman, P. S. (1988). Submarine hot springs and the origin of life. *Nature* 336:117. doi: 10.1038/336117a0
- Russell, M. J., Hall, A. J., and Martin, W. (2010). Serpentinization as a source of energy at the origin of life: serpentinization and the emergence of life. *Geobiology* 8, 355–371. doi: 10.1111/j.1472-4669.2010.00249.x
- Sabuda, M. C., Brazelton, W. J., Putman, L. I., McCollom, T. M., Hoehler, T. M., Kubo, M. D. Y., et al. (2020). A dynamic microbial sulfur cycle in a serpentinizing continental ophiolite. *Environ. Microbiol.* 22, 2329–2345. doi: 10.1111/1462-2920.15006
- Sánchez-Andrea, I., Guedes, I. A., Hornung, B., Boeren, S., Lawson, C. E., Sousa, D. Z., et al. (2020). The reductive glycine pathway allows autotrophic growth of *Desulfovibrio desulfuricans*. *Nat. Commun.* 11:5090. doi: 10.1038/s41467-020-18906-7
- Schönheit, P., Buckel, W., and Martin, W. F. (2016). On the origin of heterotrophy. *Trends Microbiol.* 24, 12–25. doi: 10.1016/j.tim.2015.10.003
- Schrenk, M. O., Brazelton, W. J., and Lang, S. Q. (2013). Serpentinization, carbon, and deep life. *Rev. Mineral. Geochem.* 75, 575–606. doi: 10.2138/rmg.2013.75.18
- Schrenk, M. O., Kelley, D. S., Bolton, S. A., and Baross, J. A. (2004). Low archaeal diversity linked to subsurface geochemical processes at the Lost City hydrothermal field, mid-Atlantic ridge. *Environ. Microbiol.* 6, 1086–1095. doi: 10.1111/j.1462-2920.2004.00650.x
- Seyler, L. M., Brazelton, W. J., McLean, C., Putman, L. I., Hyer, A., Kubo, M. D. Y., et al. (2020). Carbon assimilation strategies in ultrabasic groundwater: clues from the integrated study of a serpentinization-influenced aquifer. *mSystems* 5, e00607–e00619. doi: 10.1128/mSystems.00607-19
- Sojo, V., Herschy, B., Whicher, A., Camprubi, E., and Lane, N. (2016). The origin of life in alkaline hydrothermal vents. *Astrobiology* 16, 181–197. doi: 10.1089/ast.2015.1406
- Sumi, T., and Harada, K. (2021). Kinetics of the ancestral carbon metabolism pathways in deep-branching bacteria and archaea. *Commun. Chem.* 4:149. doi: 10.1038/s42004-021-00585-0
- Suzuki, S., Ishii, S., Hoshino, T., Rietze, A., Tenney, A., Morrill, P. L., et al. (2017). Unusual metabolic diversity of hyperalkaliphilic microbial communities associated with subterranean serpentinization at the cedars. *ISME J.* 11, 2584–2598. doi: 10.1038/ismej.2017.111
- Suzuki, S., Ishii, S., Wu, A., Cheung, A., Tenney, A., Wanger, G., et al. (2013). Microbial diversity in the cedars, an ultrabasic, ultrareducing, and low salinity serpentinizing ecosystem. *Proc. Natl. Acad. Sci. USA* 110, 15336–15341. doi: 10.1073/pnas.1302426110
- Suzuki, S., Kuenen, J. G., Schipper, K., van der Velde, S., Ishii, S., Wu, A., et al. (2014). Physiological and genomic features of highly alkaliphilic hydrogen-utilizing betaproteobacteria from a continental serpentinizing site. *Nat. Commun.* 5:3900. doi: 10.1038/ncomms4900
- Suzuki, S., Neelson, K. H., and Ishii, S. (2018). Genomic and in-situ transcriptomic characterization of the candida Monninte phylum NPL-UPL2 from highly alkaline highly reducing serpentinized groundwater. *Front. Microbiol.* 9:3141. doi: 10.3389/fmicb.2018.03141
- Tiago, I., and Verissimo, A. (2013). Microbial and functional diversity of a subterranean high pH groundwater associated to serpentinization: microbial and functional diversity of subsurface aquifer. *Environ. Microbiol.* 15, 1687–1706. doi: 10.1111/1462-2920.12034
- Truche, L., McCollom, T. M., and Martinez, I. (2020). Hydrogen and abiotic hydrocarbons: molecules that change the world. *Elements* 16, 13–18. doi: 10.2138/elements.16.1.13
- Trutschel, L. R., Chadwick, G. L., Kruger, B., Blank, J. G., Brazelton, W. J., Dart, E. R., et al. (2022). Investigation of microbial metabolisms in an extremely high pH marine-like terrestrial serpentinizing system: Ney Springs. *Sci. Total Environ.* 836:155492. doi: 10.1016/j.scitotenv.2022.155492
- Twing, K. I., Brazelton, W. J., Kubo, M. D. Y., Hyer, A. J., Cardace, D., Hoehler, T. M., et al. (2017). Serpentinization-influenced groundwater harbors extremely low diversity microbial communities adapted to high pH. *Front. Microbiol.* 8:308. doi: 10.3389/fmicb.2017.00308
- Westphal, L., Wiechmann, A., Baker, J., Minton, N. P., and Müller, V. (2018). The Rnf complex is an energy-coupled transhydrogenase essential to reversibly link cellular NADH and ferredoxin pools in the acetogen *Acetobacterium woodii*. *J. Bacteriol.* 200:e00357. doi: 10.1128/JB.00357-18
- Woycheese, K. M., Meyer-Dombard, D. R., Cardace, D., Argayosa, A. M., and Arcilla, C. A. (2015). Out of the dark: transitional subsurface-to-surface microbial diversity in a terrestrial serpentinizing seep (Manleluag, Pangasinan, the Philippines). *Front. Microbiol.* 6:44. doi: 10.3389/fmicb.2015.00044