



HAL
open science

Structural characterisation and inhibition of Arenavirus replication complex elements : assembly, function and inhibition of embedded nucleases

Sergio Hernández, Nicolas Papageorgiou, Maria Spiliopoulou, Mikael Feracci, Laura Garlatti, Barbara Selisko, Afroditi Vaitsopoulou, Thi-Hong van Nguyen, Clemence Mondielli, Magali Saez-Ayala, et al.

► To cite this version:

Sergio Hernández, Nicolas Papageorgiou, Maria Spiliopoulou, Mikael Feracci, Laura Garlatti, et al.. Structural characterisation and inhibition of Arenavirus replication complex elements : assembly, function and inhibition of embedded nucleases. 26TH CONGRESS AND GENERAL ASSEMBLY OF THE INTERNATIONAL UNION OF CRYSTALLOGRAPHY, Aug 2023, Melbourne (AUS), Australia. hal-04180334

HAL Id: hal-04180334

<https://amu.hal.science/hal-04180334v1>

Submitted on 11 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Structural characterisation and inhibition of Arenavirus replication complex elements : assembly, function and inhibition of embedded nucleases.

S. Hernandez¹, N. Papageorgiou¹, M. Spiliopoulou³, M. Ferracci¹, L. Garlatti¹, B. Selisko¹, A. Vaitsooulou¹, T-H. V. Nguyen¹, C. Mondielli¹, M. Saez-Ayala¹, Y. Laban¹, I. Margiolaki³, B. Canard^{1,2}, K. Alvarez¹, **F. Ferron**^{1,2*}

¹ Architecture et Fonction des Macromolécules Biologiques - CNRS - Aix-Marseille Université - UMR7257 - Viral Replicases: Structure, Mechanism, and Drug-Design, 13009, Marseille, France.

² European Virus Bioinformatics Center, Leutrargraben 1,07743 Jena, Germany.


³ Section of Genetics, Cell Biology and Development, Department of Biology, University of Patras, GR- 26500 Patras, Greece.

*Correspondence to : francois.ferron@univ-amu.fr


A007 / 530 P0019

INTRODUCTION

Arenaviruses, belongs to a family of emerging enveloped segmented and ambisens RNA viruses associated with neurological and hemorrhagic diseases in humans. Arenavirus transcription and genome replication are cytoplasmic ensured by a ribonucleoprotein replicase complex NP-L. After penetration, L protein initiates transcription to produce NP and L mRNAs [1]. The priming of transcription is the result of a *cap-snatching mechanism* ensured by an endonuclease domain associated to the L polymerase. As the concentration of NP in the cell increases, genome segments are replicated, to produce full-length copies (cRNA). cRNAs are now templates for transcription of GPC mRNA (from the S segment) and Z mRNA (from the L segment). The NP carries an *exonuclease* in charge of clearing out from the cytoplasm dsRNA triggering innate immunity response. Both nucleases have a similar *two metal ion catalytic mechanism*, with the particularity of transitioning ion brought by the RNA substrate. Any alteration of the remaining ion impairs greatly these activities[2]. We present a global study aiming to characterize the assembly of the NP[3], through flexible domains[4], a step critical for vRNA packaging and the positioning of L for vRNA replication, as well as using a combined approach of biophysical screening, *crystallography* and *in silico* docking, identifying active compounds against both nucleases[5]. Crystal structures of the nucleases domain complexed with several compounds were obtained[6,7]. By developing specific compounds to alter both transcription and innate immunity shadowing, our strategy is to give the cell a fighting chance to clear the infection. Combining structure, enzymology, rational synthesis, hit-to-lead optimization, *in cellula* evaluation, and screening methods, we are presenting the results of a 2nd generation of molecules paving the way to the design of a 3rd generation increasing specificity towards Arenaviral nucleases in the context of the replication complex[8].




STRATEGY




RESULTS


NP characterization




Structural model fitting structural data




Structural similarities with *Nairovirus*' N-ExoN MOA & Inhibition by ion chelation




ExoN MOA & Inhibition by ion chelation




Arenavirus pipe-line




FP signal recorded during 30 min, every 30s. RNA substrate concentration (100 nM). ExoN concentration (0.1 μM to 1.6 μM). Average ± SEM of three independent experiments.




Scatter plot showing the % of inhibition



Screening Methodology



Arenavirus RNP Reconstruction



CONCLUSIONS & PERSPECTIVES

Global comprehension of biological system is critical to achieve proper understanding of the MOA of an enzyme and to develop inhibition strategy. Over the last decade we have characterized several structures and activities of the key proteins of Arenaviral replication complex, both apo and in complex with inhibitors targeting the two nucleases. Compounds targeting both viral transcription and viral impairment of innate immunity present lesser the chances of selecting resistant mutations. We proved thru transversal approach studies that fluorescence polarization (FP) method is reliable, sensitive to monitor nuclease activity and feasible in 384-well plates. The method is robust, rapid and non-destructive allowing the visualization of the RNA substrate degradation using gels as orthogonal method of confirmation. We identified several hit inhibitors of viral arenavirus (LCMV) ExoN activity and determined by fluorescence polarization their IC₅₀ at a micromolar level. Then we validated their activities in cell culture assays and we found compounds displaying efficiency at micromolar level against Junin virus a new world Arenavirus. These results highlight the effectiveness of translational approach in the screening of compounds libraries for the discovery of antivirals. Structural characterization of the compound in complex with the ExoN is underway.

REFERENCES

[1] Ferron F, Weber F, de la Torre JC, Reguera J. *Virus Res.* 2017 Apr;15:234.
 [2] Yekwa E, et al. *Acta Crystallogr D Struct Biol.* 2017 Aug;17(8):641-649
 [3] Papageorgiou N, et al. *F. FEBS Open Bio.* 2021 Apr;11(4):1076-1083.
 [4] Papageorgiou N, et al. *Viruses.* 2020 Jul;12(7):772
 [5] Hernández S, et al. *Antiviral Res.* 2022 Aug;204:105364
 [6] Saez-Ayala M, et al. *IUCr.* 2018 Feb;22(5):223-235.
 [7] Nguyen THV, et al. *IUCr.* 2022 May;28(9):468-479
 [8] Saez-Ayala M, et al. *Antiviral Res.* 2019 Feb;162:79-89.

Acknowledgements

