



HAL
open science

Optimized miR-124 reporters uncover differences in miR-124 expression among neuronal populations in vitro

Catherine Lepolard, Cynthia Rombaut, Florence Jaouen, Ana Borges, Elodie Caccamo-Garcia, Natalia Popa, Eduardo Gascon

► To cite this version:

Catherine Lepolard, Cynthia Rombaut, Florence Jaouen, Ana Borges, Elodie Caccamo-Garcia, et al.. Optimized miR-124 reporters uncover differences in miR-124 expression among neuronal populations in vitro. *Frontiers in Neuroscience*, In press. hal-04221595

HAL Id: hal-04221595

<https://amu.hal.science/hal-04221595v1>

Submitted on 29 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Title: Optimized miR-124 reporters uncover differences in miR-124 expression among neuronal populations in vitro

Authors: Catherine Lepolard^{1,†}, Cynthia Rombaut^{1,†}, Florence Jaouen^{1,2,†}, Ana Borges¹, Elodie Caccomo-Garcia¹, Natalia Popa¹, and Eduardo Gascon^{1,*}

¹ Aix Marseille Univ, CNRS, INT, Inst Neurosci Timone, Marseille, France

² NeuroBioTools facility (NeuroVir), Aix Marseille Univ, CNRS, INT, Inst Neurosci Timone, Marseille, France

† These authors share first authorship

*Corresponding author: Eduardo Gascon, Institut de Neurosciences de la Timone, Aix-Marseille Université; CNRS UMR7289; 27, Boulevard Jean Moulin, 13005 Marseille (France); E-mail: eduardo.gascon-gonzalo@univ-amu.fr

Running title: Optimized miRNAs reporters

ABSTRACT

Although intensively studied in the last decades, how microRNAs (miRNAs) are expressed across different cell types in the brain remains largely unknown. To address this issue, we sought to develop optimized fluorescence reporters that could be expressed in precise cellular subsets and used to accurately quantify miR contents in vivo. Focusing on miR-124, we tested different reporter designs whose efficiency was confirmed in different in vitro settings including cell lines and primary neuronal cultures from different brain structures. Unlike previous reporters, we provide experimental evidence that our optimized designs can faithfully translate miR levels in vitro. Such tools would enable assessing miRNA expression at the single cell resolution and are expected to significantly contribute to future miRNA research in vivo.

Keywords: miRNA, miR-124 reporters, neuron, AAV, mice

INTRODUCTION

microRNAs (miRNAs) are a class of short (20-25 nt) non-coding RNAs (Huntzinger and Izaurralde, 2011; Izaurralde, 2015). miRNAs repress expression of messenger RNAs (mRNAs) containing short complementary sequences. Since these sequences could be found in hundreds of transcripts, miRNAs are thought to fine-tune gene expression at a global scale (Fabian and Sonenberg, 2012). Mechanistically, it has long been known that miRNA-dependent silencing critically relies on the complementarity of miRNA-mRNA sequences. The complementarity in first eight nucleotides of the miRNAs (the so-called seed region) is critical for target recognition whereas mismatches in the remaining sequence are tolerated and do not seem to affect miRNA binding efficiency (Huntzinger and Izaurralde, 2011). Nonetheless, perfect base-pairing has been consistently shown to result in transcript degradation whilst translational inhibition is the main silencing mechanisms for those targets with lower degree of complementarity (Valencia-Sanchez et al., 2006; Yekta et al., 2004).

miRs have attracted much attention in the context of neurodegenerative diseases. On one hand, because a growing body of evidence suggest that miRNAs dysregulation is a key pathogenic event in multiple diseases (Gascon et al., 2014; Liu et al., 2012; Salta and De Strooper, 2017). On the other hand, because of their simultaneous action on different biological pathways, they hold a promising potential for therapeutic applications (Gentile et al., 2022; Seyedaghamiri et al., 2023).

miR-124 is one of the best studied miRNAs because of: i) its abundance (Lagos-Quintana et al., 2002) ;ii) conservation across species and brain enrichment (Lagos-Quintana et al., 2002); iii) involvement in multiple key biological processes in the brain ranging from neuronal differentiation during development (Maiorano and Mallamaci,

2010; Makeyev et al., 2007) to fine-tuning neurotransmitter receptor composition at the synapse (Gilbert et al., 2016; Hou et al., 2015; Namkung et al., 2023); and iv) alterations observed in a wide variety of pathological conditions, especially neurodegenerative diseases (Gascon et al., 2014; Ghafouri-Fard et al., 2022; Wang et al., 2018; Yao et al., 2019). Remarkably, there is currently no information about miR-124 expression at the single-cell resolution.

Owing to their small sizes and relative low abundance, miRNA quantification has been technically challenging. miRNA expression has been evaluated using in situ hybridization (Urbanek et al., 2015), northern blot (Kim et al., 2010), microarrays (Cissell and Deo, 2009), RT-PCR amplification (Chen et al., 2005) and, more recently, sequencing (small RNA sequencing) (Podnar et al., 2014; Zhou et al., 2011). However, except last-generation in situ hybridization with RNAscope plus probes, precise quantification and cellular resolution remains an unsolved problem. The advent of single-cell RNA sequencing (scRNA-seq) technology has enabled transcriptomic profiling of individual cells and enormously contributed to our understanding of brain complexity at the molecular level. Nevertheless, current scRNA-Seq methods are restricted to long RNAs, primarily mRNAs (Macosko et al., 2015; Rosenberg et al., 2018). In summary, due to technical constraints, how miRNAs are expressed in different cell types in vivo remains elusive.

To monitor miRNA levels, several groups have devised different reporter strategies (reviewed in (Song et al., 2020)). Among them, miRNA binding sequence-engineered fluorescent reporters have been particularly attractive, as they might enable both precise quantification and visualization of miRNA level in living cells (Kato et al., 2010;

Sano et al., 2017; Wang et al., 2019). In the same line, several transgenic reporter mice strains have been generated aiming at characterizing the cellular expression of different miRNAs in the brain in vivo (Åkerblom et al., 2014; Åkerblom et al., 2012; Åkerblom et al., 2013). Previous miRNA reporter constructs have been engineered to contain multiple (4 or more) perfectly complementary sequences for a particular miRNA. Such reporters are submitted to a strong inhibition in vivo so that only cells completely devoid of the miRNA show reporter expression (Keaveney et al., 2018; Keaveney et al., 2020; Åkerblom et al., 2012; Åkerblom et al., 2013). These observations argue for a non-physiological regulation of this kind reporter. In summary, new reporters are therefore required to circumvent the above-mentioned technical hurdles.

Here, we reasoned that mimicking endogenous targets should be more appropriate for research purposes. We sought to develop reporter designs based on more physiological binding sequences for, We choose miR-124 as a prototypical miRNA involved in multiple brain diseases. Setting up such miR-124 reporters could not only provide important insights into the generalization of our approach to other miRNAs but also, because of its conservation, it could serve to interrogate different species. Here, we investigated in vitro the activity of our novel reporters and compared them with previous designs. Using different cell lines and primary neurons, our results show that reporters bearing endogenous binding sequences are more reliable to evaluate miR-124 expression. Overall, we provided novel tools that can be useful for the investigation of expression profiles of specific miRNAs at the cellular resolution and, more importantly, that should be applicable to monitor/identify specific cell populations in vivo.

METHODS

miR-124 reporter design

For ETS design, we selected the binding sequence found in the mouse *Gria2* mRNA (at position 172-192 in the UTR). This choice was made based on our previous experience (Gascon et al., 2014) as well as the observations from other groups (Dutta et al., 2013; Namkung et al., 2023) having demonstrated that *Gria2* is a relevant target for miR-124 in rodents and humans. For PCS, we used the complementary miR-124 sequences. For mutated versions, we replaced three nucleotides of the binding sequence from CUU to GGG. These nucleotides base pair within the seed region (position 2-4) and seven. All sequences were shown in Table 1 as well as in Suppl Fig 1a. All sequences were cloned into the 3' UTR of a nuclear RFP (H2B-RFP) for easy quantification using cell cytometry.

Cloning of binding sequences

Perfectly complementary sequences or endogenous sequences were cloned into a plasmid backbone containing a H2B-RFP using appropriate restriction enzymes. For miR-124 overexpression, 2 kb of the primary miR-124 were amplified from mouse genomic DNA using PCR and then cloned into a plasmid allowing the co-expression of H2B-GFP. All plasmids used here contained Tol2 sequences for the generation of stable cell lines (see below). The constructs encompassing the H2B-RFP and the different 3'UTR were amplified using PCR and cloned in an inverted orientation into a pAAV-DIO vector (Addgene #35507). Plasmids maps are shown in Suppl. Fig. 4.

Viral production

Conditional reporters AAVs were generated using pAAV-DIO plasmids described before. Synapsin-Cre AAVs were produced from Addgene plasmid #105540. Viral productions were carried out using standard protocols with slight modifications (Su et al., 2020). Briefly, HEK293T cells were transfected using the calcium phosphate method (pHelper, pRC2, pAAV; ratio 2:1:1). After 60-72h, cells were harvested and AAV particles purified and concentrated using Takara dedicated kit (Takara, France). Viral productions were tittered using quantitative PCR (Takara, France). AAV9 serotypes of a titer around 10^{12} viral genomes/ml were used in this report. The different vectors are listed in Table 2.

Construction of stable cell lines

HEK293T cells were obtained from ATCC and cultured under standard conditions (DMEM-10% Fetal Calf serum). They were used to generate cell lines stably expressing miR-124 as well as the different reporters. For that, we used a previously described procedure (Kawakami and Noda, 2004). Briefly, the different constructs were cloned into a vector containing the Tol2 sequences for the medaka fish transposase. Then, these plasmids were co-transfected with a plasmid containing the transposase into HEK cells. After a clonal dilution, cells having stably incorporated the transgene were selected by fluorescence expression. The expression of the proper construct was verified using quantitative PCR.

Primary neuronal cultures

Cell cultures were prepared from newborn CD1 mice (postnatal day 0). Briefly, animals were decapitated, the brains rapidly transferred into ice-cold Hank's balanced salt solution (HBSS, Gibco, France) and the different brain dissected out.

Then, they are dissociated mechanically, trypsinized and purified using percoll gradient centrifugation. Cells were plated onto poly-ornithine (Sigma, France) coated 6-wells plates and allowed to grow in Neurobasal medium (Gibco) with 2% B27 supplement (Gibco), 2 mM glutamate (Gibco), 1 mM sodium pyruvate (Gibco) and antibiotics (Gibco). Cell density at plating was 3×10^5 cells/well Half of the medium was replaced twice a week.

Transfection and transduction

In all experiments with cell lines, we used Lipofectamine 3000 (Thermo) as transfection reagent following manufacturer's recommendations. For primary cultures, AAVs particles (2 μ l/well, 1 μ l from the reporter AAV and 1 from the Cre AAV, representing around 10^{10} genomes/culture) were diluted in 50 μ l of culture medium and then added to the cells one day after plating (DIV1). Neurons were analyzed 10 days after viral infection.

Cell cytometry and FACS sorting

Cells were gently detached using trypsin and rinsed in PBS. After removal of cell clumps (70 μ m cell strainer), individual cells were resuspended into 500 μ l of PBS-0.1% BSA. The cells were analyzed on a cytoFlex device(Beckman Coulter) and data were analyzed with the CytoExpert software (Beckman Coulter). For FACS, infected neurons were sorted using a MoFlo Astrios EQ device (Beckman Coulter). Negative cells (either not transfected or transduced) were used to set up the gates.

miRNA quantification

Total RNA containing miRNAs and mRNAs was then extracted from these samples using NucleoSpin miRNA kit (Machery Nagel) following manufacturer's instructions. RNA extraction protocol includes an on-column DNase treatment (30 min). Total RNA was quantified using a NanoDrop (Fisher Scientific).

For miRNA quantification, we first performed reverse transcription using TaqMan advanced miRNA cDNA synthesis kit (Applied Biosystems, France) according to the protocol provided by the manufacturer. Abundance of miRNAs in the samples was measured using custom TaqMan probes (Applied Biosystems, USA) and a QuantStudio 7 PCR thermocycler (Applied Biosystems, France). Relative quantification was carried using the $\Delta\Delta\text{Ct}$ method. Reference genes were GAPDH (Fig. 1c), miR-92 (Suppl Fig 2), miR-16 and miR-101 (Fig 3a) and miR-9 (Fig 3d). For quantification of miR-124 in FACS sorted cerebellar neurons, we did not use miR-16 or miR-101 because their expression is much lower than miR-124 and could not be consistently amplified in these samples. miR-9, a neuronal enriched miRNA, was preferred as it was the miRNA whose levels were closer to miR-124 in these cultures. All qPCR assays (including for AAV titration) are commercially available and their references indicated in Table 3.

Statistics

Because of the limited sample size in our experiments (n=3 or 4), we could not perform normality checks. To avoid using low-power non-parametric tests (i.e. Kruskal Wallis), we transformed our data using a relative normalization method. Instead of comparing the absolute fluorescence intensity, we calculated the percentage of fluorescence in each experiment taking the RFP control sample as reference (whose intensity was arbitrarily set to 100). For the detection of

significant differences among groups, we used therefore ANOVA analysis followed by post-hoc tests. Data is presented as means \pm SEM unless indicated. All statistical analysis was performed using Prism GraphPad software (version 9). Details of tests and n for each experiment are provided in Figure legends.

RESULTS

miR-124 reporter designs based on endogenous binding sites

We generated and systematically tested in vitro different fluorescent reporter designs for miR-124. Our constructs encompass a nuclear RFP and different numbers (1 or 2) and types (perfectly complementary or sequences found in endogenous targets) of miR-124 target sequences (Fig 1a and Suppl Fig 4). We first generated stable cell lines containing such reporters and evaluated how miR-124 expression influenced their activity. For that, we transiently transfected these different lines with a plasmid driving the expression of miR-124 and GFP and use untransfected cells as a reference for RFP intensity. As expected, cells transfected with this plasmid, we did not observe any change in RFP fluorescence in GFP⁻ cells (Suppl. Fig 1a). A significant reduction of RFP fluorescence was already observed 2 days after transfection in cells expressing the perfectly complementary sequences (PCS). Silencing was exacerbated at 6 days post-transfection leading to a near complete extinction of fluorescence and confirming previous reports. The effect was similar in cells containing either one or two PCS.

In sharp contrast, reporters bearing endogenous sequences (ETS) exhibit a more modest decreases in fluorescence and a slightly additive effect of increasing the number of target sequences (Fig 1b). This moderate effect might reflect a regulation comparable to that of endogenous targets.

To further understand the effect of miR-124 on the different reporters, we FACS sorted transfected cells and analyzed reporter expression at the mRNA level. Our

analysis (Fig 1c and Suppl Fig 1b) revealed striking differences of miR-124 effect on the reporters bearing ETS and PCS. Levels of RFP transcript were similar in GFP⁻ and GFP⁺ cells in the lines containing ETS reporters. Conversely, a significant reduction of RFP mRNA was found in lines bearing PCS. Together with our observations on fluorescence intensity, those results suggest that translation is inhibited for ETS reporters whereas both transcript degradation and translation blocking operate to silence PCS reporters. Interestingly, despite the fact that miR-124 levels in GFP⁺ cells increased along with time (Suppl Fig 1c), there is no change in the abundance of RFP transcripts neither in ETS nor in PCS-bearing reporters from 2 to 6 days indicating a different time course between mRNA and protein regulation. These results indicate that ETS and PCS can be useful to estimate miR-124 expression at the protein and the mRNA level, respectively.

miR-124 reporters bearing endogenous target sequence enable discrimination of HEK cells engineered to express different miR-124 contents

Since the final aim of reporters is uncovering different levels of miR-124 across different cell types in vivo, we explored whether our different designs can detect different miR-124 contents in vitro. We engineered HEK cells, endogenously devoid of miR-124 (Gascon et al., 2014; Visvanathan et al., 2007; Xue et al., 2016), to stably express different amounts of this miRNA . We selected two cell lines displaying moderate but different levels of miR-124 to approach more physiological conditions (Suppl Fig 2a).

We then analyzed the reporter activity of our different constructs in these cell lines. To avoid the intrinsic variability associated with transient tranfection (in terms of

copy number in each cell and plasmid dilution), we constructed dual reporter plasmids (Suppl. Fig 4) containing not only a miR-124 reporter but also a reference reporter (CFP) that serves as internal reference (Suppl Fig 4). Using flow cytometry, we quantified the intensity of both RFP and CFP fluorescence. The RFP/CFP ratio was calculated as a reliable output of miR-124 effects on the RFP reporter. We observe that PCS containing reporters are subjected to strong inhibition (Fig 2b). As before, we found an almost identical silencing independently of the cell line and the number of binding sequences. This suggest that PCS reporter are extremely sensitive even to moderate miR-124 levels. ETS-containing reporters, in turn, exhibit a more tunable regulation and a silencing proportional to the number of sequences present in the reporter (Fig 2c). As a control, transfection of all these constructs in regular HEK cells resulted in no modification of RFP/CFP ratio (Suppl Fig 2). Additionally, mutating 3 nucleotides of the seed region in 1x ETS reporter abrogate the regulatory effect of miR-124 arguing for the specificity of our reporters (Suppl Fig 2). These observations together indicate that reporters containing ETS might be a more appropriate strategy to measure miR-124 activity in vivo.

Optimized miR-124 reporter translates miR-124 levels in cultured neurons

We next tested one of our ETS designs in a more physiological setting, primary neuronal cultures from the mouse hippocampus, cortex and cerebellum. We selected them because, as shown in Fig 3a, cultured neurons from these brain regions express different endogenous levels of miR-124 but not of other miRNAs.

To introduce different reporters in primary neurons, we used adeno-associated vectors (AAVs) as previous work have shown their efficiency and lack of toxicity (Malik et al., 2012). We constructed three Flex conditional AAVs containing

respectively a control reporter (no miR-124 binding sequence, AAV-RFP), a reporter bearing a single ETS (AAV-RFP-miR-124) and a mutated version (3 nucleotide mismatch in the seed region) of the reporter (AAV-RFP-miR-124mut). We transduced different neuronal cultures with the reporter AAVs along with a synapsin-Cre-GFP AAV. This strategy enables not only to drive the reporter express in similar neuronal subsets in the cultures but also to have a reference fluorescent protein from the Cre virus. We observe that AAVs transduce primary cultures with a high efficiency (ranging from 30-60% of hippocampal cells, 15-30 % of cortical neurons and 50-75% of cerebellar culture, Suppl Fig 3a). As expected, the GFP fluorescence and the proportion of GFP+ cells does not differ among the different conditions confirming that reporter proteins are not subjected to any unspecific silencing (Suppl Fig 3a and b). In contrast, compared to the control AAV, neurons transduced with miR-124 reporter show a marked reduction in the proportion of RFP+ cells (Suppl Fig 3a) as well as in the intensity fluorescence (Suppl Fig 3c). The intensity of such decrease varies among the brain structures analyzed. Thus, cerebellar neurons exhibit the highest silencing and hippocampal cultures are less inhibited (Fig 3b). Mutation in the miR-124 binding sequence abolishes such effect arguing for the specificity of our results. To further support these observations, we also calculated the RFP/GFP ration as a more robust quantitative measure of reporter fluorescence (Fig 3b). In the same line, we detected significant changes in neurons trasnduce with the miR-124 reporter.

To confirm these observations and to avoid the variability among different experiments, we sought to analyze the activity of our reporters in neurons from the same culture. We reasoned that, if our reporter fluorescence is negatively correlated to miR-124 contents, neurons displaying high RFP levels (RFP^{high}) might contain

lower amounts of miR-124 relative to those in which RFP silencing is stronger (RFP^{low}). Conversely, in neurons transduced with the control or mutated reporter, miR-124 levels should be no correlated to fluorescence intensity. We therefore transduced cerebellar neurons (those showing highest miR-124 expression and the highest transduction efficiency), carried out FACS sorting to isolate RFP^{high} and RFP^{low} neurons and measured miR-124 in these different populations. We first confirmed that, in neurons transduced with the miR-124 reporter, there was an obvious left-shift in the fluorescence distribution that resulted in a net decrease of RFP^{high} cells (Suppl Fig 3c). Regarding miR-124 contents, neurons having received the control or mutated construct show similar miR-124 contents irrespective of the fluorescence intensity (Fig 3c). In contrast, in those transduced with miR-124 reporter, RFP^{low} neurons exhibit a significant enrichment in miR-124 levels compared to RFP^{high} neurons suggesting that our reporter is subjected to miRNA silencing and its levels inversely correlated to miR-124 (Fig 3c). Finally, using linear regression, we sought to confirm the inverse correlation of reporter fluorescence and miR-124 contents in FACS-sorted cerebellar neurons. For that, we plotted the mean RFP fluorescence and relative miR-124 levels in each condition. As shown in Fig 3d, linear correlation was remarkably good for the miR-124 reporter ($R^2=0.6584$) but rather poor for the control RFP reporter as well as in the mutated reporter ($R^2=0.01877$ and $R^2=0.004265$, respectively). More importantly, the slope analysis revealed a significant difference from 0 only for miR-124 reporter ($p=0.01255$) but not the control reporters ($p=0.8955$ and 0.9022). These findings provide further support for the inverse relationship between miR-124 abundance and reporter fluorescence. Overall, our in vitro experiments support the notion that optimized miR-124 reporters could be useful for in vivo applications.

DISCUSSION

In this work, we optimized an existing technique (fluorescent reporters) to infer miR-124 levels in vitro. Our approach relies on the expression of a nuclear reporter fluorescent protein bearing endogenous binding sequences for miR-124. Our reporters showed a moderate, more physiological silencing which enables the detection of miR-124 differences among distinct neuronal types in culture. This is in sharp contrast to recent studies in which miR-124 reporters (or reporters for other miRNAs) are submitted to strong inhibition (Åkerblom et al., 2014; Åkerblom et al., 2012; Åkerblom et al., 2013). Our results show an important difference in the mechanism of silencing among sequences; PCS destabilizes the transcript and block protein translation whereas ETS do only act at the translational level. These results are in line with previous work indicating that endonucleolytic cleavage is favored by perfect base-pairing between the miRNA and the mRNA (Valencia-Sanchez et al., 2006; Yekta et al., 2004) and might explain the striking different levels of inhibition across reporters bearing PCS and ETS sequences.

Although intensively studied, our understanding of miR-124 expression levels across cell types remains rudimentary. The main aim of this study is to generate miR-124 reporters that could help unraveling the precise expression pattern of this miRNA in the brain. More precisely, miR-124 reporters are intended to: i) estimate the abundance of this miRNA in living cells and ii) compare expression among different neuronal subsets. Concerning the first point, this is a key aspect as most techniques aiming at measuring and comparing miRNAs expression levels are end-point techniques (quantitative PCR, miRNA sequencing, in situ hybridization or miRNA

microarrays). This is precluding the prospective identification of specific neuronal subtypes according to the expression of a particular miRNA and the further investigation of how changes in this miRNA alter neuronal properties in different cellular contexts. Regarding the expression of a particular miRNA in precise cell types, we have provided evidences that miRNA reporters could be packed in Cre-dependent AAVs and delivered to specific cell subsets so that, in combination with transgenic Cre lines, they can be used to thoroughly map the expression of miR-124 during development, aging or in pathological conditions.

In this work, we provided initial evidence that reporters bearing ETS could be more appropriate to estimate miR-124 levels *in vitro*. The *in vivo* application of approaches described here is more challenging. These experiments rely on the stereotactic injection of the reporter AAV into a target brain area. This approach introduces important bias in the number of transduction units delivered to each cell as the number of viral particles exponentially decrease with the distance. Thus, those cells near the injection site would receive high number of particles potentially driving reporter levels out of the regulatory range of miR-124 *in vivo*. Conversely, those far from the injection might receive suboptimal number of AAVs particles so that reporter expression after miR-124 silencing does not reach detection thresholds. To circumvent such limitations, *in vivo* applications would require an optimization of viral constructs: i) promoter choice so that the reporter levels are appropriate independently of the number of AAV copies in each cell; ii) including a second reporter not subjected to miR-124 silencing so that it can be used as internal reference for AAV infection. *In vivo* validation of miR-124 reporters will be a focus of our future work.

Research on miRNAs has been exponential in the last two decades. This interest has been boosted, in part, by the paradigm shift in pharmacological approaches against complex human disorders such as neurodegenerative diseases (Lee et al., 2021; Millan, 2017; Nguyen et al., 2022; Wang et al., 2018). Rather than looking for one target, current therapeutic strategies aim at targeting multiple pathways shared by a number of conditions (Grosjean et al., 2022; Ibrahim and Gabr, 2019). miRNA-based therapy is particularly suited for such purposes, as their mechanism of action is grounded on regulation of multiple components across molecular cascades that might converge on disease-relevant alterations (Kosik and Krichevsky, 2005). Although promising, application of miRNAs to the clinic requires further fundamental research to better characterize not only how miRNAs target key pathological pathways but, more importantly, which are the cells types where they should be delivered. New tools are required not only to increase our understanding of miRNA expression patterns in the brain but also to monitor cell-specific changes in miR-124 levels in animal models of neurodegenerative disorders (Tran et al., 2023; Zhao et al., 2023). Our work opens a potential new option for the design of such tools.

A major limitation of our study concerns the reporter design. Since miRNA binding silence the expression of the target mRNA, our reporter is inversely correlated to the actual miRNA levels. A reporter showing a direct relationship to the miRNA contents might be preferable in certain contexts. In addition, it remains to be determined whether reporter bearing one target endogenous sequence are functional for other miRNAs. Despite these limitations, reporters tested here should provide a rationale

for future designs and hold a great potential to unravel miR-124 expression in the brain.

Data Availability Statement

All data supporting the conclusions are included in the article/supplementary material. Further inquiries can be directed to the corresponding author.

Author Contributions

CL and CR performed most of the experiments and helped out in the writing of the first draft of the manuscript. FJ: viral strategy and production, writing the manuscript; AB and ECG: provided essential help on early stages of the project; NP cytometry analysis. EG: writing, review and editing, supervision, project administration and funding acquisition. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by the French National Agency (ANR) to EG (ANR-22-CE17-0034), Fondation de France (00100077), and Fondation France Alzheimer (2021-#6239).


Conflict of Interest

The authors report no biomedical financial interests or potential conflicts of interest.


Acknowledgements

The authors would like to thank Thomas Bescond for the viral productions, Stephane Robert from AMUTICYT facility for technical support with cytometry and FACS sorting and Dipankar Bachar for his feedback on bioinformatics/statistical analysis.

a



b



c




Figure legends

Figure 1. Testing reporter activity in cell lines engineered to express different miR-124 reporter constructs.

- a) Schematic representation of experimental protocol followed in these experiments (upper panel) as well as the different miR-124 reporter designs tested (lower panels)
- b) Effect of miR-124 transfection on RFP fluorescence in cells lines stably expressing a control reporter (RFP with no binding sequence for miR-124), PCS-containing (left panel) or ETS-containing miR-124 reporter (right panel). Cytometry analysis confirmed a reduction of RFP fluorescence in all miR-124 reporters. PCS showed a much more profound silencing effect than ETS. No significant changes were found in cells bearing 1 or 2 binding sequences. Two-way ANOVA, Tukey post-hoc test (n=3 independent transfections of the different cell lines).
- c) Effect of miR-124 transfection on RFP mRNA levels. 2 and 6 days after transient transfection with a miR-124-GFP plasmid, GFP⁺ cells were FACS sorted and the levels of RFP transcripts were measured using quantitative RT-PCR. Using GAPDH as reference gene, we observed a significant reduction of RFP mRNA only in PCS-containing miR-124 lines. Two-way ANOVA, Holm-Sidak post-hoc test (n=3 independent transfections of the different cell lines).

* p<0.05 ** p<0.01 *** p<0.001






Figure 2. Testing miR-124 reporter designs in cell lines engineered to express different miR-124 levels.

- a) Schematic representation of experimental protocol followed in these experiments.
- b) Cells expressing low and moderate levels of miR-124 were transiently transfected with PCS-containing miR-124 RFP reporters and a CFP control plasmid. Cytometry analysis revealed a strong reduction of fluorescence ratio compared to cells transfected with a control confirmed a reduction of RFP fluorescence. Two-way ANOVA, Tukey post-hoc test (n=3 independent transfections of the different cell lines).
- c) Similar experiments were performed using ETS-containing miR-124 RFP reporters as well as the CFP control plasmid. These constructs show a less intense but still significant reduction in fluorescence ratio. Two-way ANOVA, Tukey post-hoc test (n=3 independent transfections of the different cell lines).

* p<0.05 ** p<0.01 *** p<0.001

,




Figure 3. Testing reporter activity in primary neuronal cultures from cortex, cerebellum and hippocampus.

- a) Levels of different neuronal enriched miRNAs in cortical, hippocampal and cerebellar cultures (n=5 for cortical and hippocampal cultures, n=10 for cerebellar cultures).
- b) Neuronal cultures were transduced with AAVs driving the expression of different floxed-RFP reporters as well as AAV-Syn-Cre-GFP. 10 days after viral infection, intensity of the RFP (from the reporter AAV) and GFP (from Syn-Cre AAV) were measured and their ratio calculated. miR-124 reporter show a significant reduction of the ratio in all neuronal cultures but particularly in the cerebellum. Mutation of 3 nucleotides in the seed region (nt 2-4) abrogated this effect. Two-way ANOVA, Dunnett post-hoc test (n=3 independent for hippocampus and cortex, n=4 for cerebellum).
- c) Analysis of miR-124 expression in cerebellar cultures transduced with AAV reporters. Using FACS, we sorted primary cerebellar neurons and quantified miR-124 expression in RFP^{low} and RFP^{high} cells taking RFP⁻ cells as reference. We showed that miR-124 levels were significantly lower in RFP^{high} neurons in cultures transduced with the AAV-RFP-miR-124. This effect can not be observed in cultures transduced with the control AAVs (bearing a control RFP or a mutated 1x ETS reporter). Two-way ANOVA. Bonferroni post-hoc test (n=4 independent for AAV-RFP and AAV-miR-124, n=3 for AAV-miR-124mut).
- d) Linear regression analysis of RFP fluorescence and miR-124 contents in cerebellar FACS-sorted neurons. When we plotted the mean fluorescence of RFP and the relative miR-124 levels, we only observed a linear relationship ($R^2=6584$) and a slope different from 0 ($p=0.0145$) in neurons transduced with the miR-124 reporter AAV (n=4 independent for AAV-RFP and AAV-miR-124, n=3 for AAV-miR-124mut).

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

Table 1. Sequences used for ETS and PCS binding sequences

Binding sequence	Nucleotide sequence
1x PCS	GGCATTACCGCGTGCCTTA
2x PCS	GGCATTACCGCGTGCCTTAATTCGAATGGCATTACCGCGTGCCTTA
1x ETS	GGAACCTTCTGAGTGCCTTA
2x ETS	GGAACCTTCTGAGTGCCTTAATTCGAATGGAACCTTCTGAGTGCCTTA
1x ETS mut	GGAACCTTCTGAGTGCGGGA

Table 2. Viral vectors used in this study

Name	Serotype	Titer (genomes/ml)
Syn-Cre-GFP (generated from Addgene plasmid #105540)	AAV9	$2,4 \times 10^{12}$
RFP (no binding sequences for miR-124)	AAV9	$1,3 \times 10^{12}$
miR-124 (1x ETS for miR-124)	AAV9	$9,5 \times 10^{11}$
miR-124mut (1x ETS for miR-124 mutated in 3 nucleotides pos 2-4)	AAV9	3×10^{12}

Table 3. Quantitative PCR assays used in this study

Name	Manufacturer	Assay ID
miR-124-3p	Thermo	mmu480906_mir
miR-9-5p	Thermo	mmu481285_mir
miR-125a-5p	Thermo	mmu480906_mir
miR-92	Thermo	477827_mir

Name	Manufacturer	Assay ID
RFP	Thermo	Mr07319438_mr
GAPDH	Thermo	Hs99999905_m1

Name	Manufacturer	Catalog #
AAVpro Titration Kit	Takara	6233

REFERENCES

- Akerblom, M., Petri, R., Sachdeva, R., Klussendorf, T., Mattsson, B., Gentner, B., & Jakobsson, J. (2014). microRNA-125 distinguishes developmentally generated and adult-born olfactory bulb interneurons. *Development*. 141, 1580-1588.
- Åkerblom, M., Sachdeva, R., Barde, I., Verp, S., Gentner, B., Trono, D., & Jakobsson, J. (2012). MicroRNA-124 is a subventricular zone neuronal fate determinant. *J Neurosci*. 32, 8879-8889.
- Åkerblom, M., Sachdeva, R., Quintino, L., Wettergren, E. E., Chapman, K. Z., Manfre, G., Lindvall, O., Lundberg, C., & Jakobsson, J. (2013). Visualization and genetic modification of resident brain microglia using lentiviral vectors regulated by microRNA-9. *Nat Commun*. 4, 1770.
- Chen, C., Ridzon, D. A., Broomer, A. J., Zhou, Z., Lee, D. H., Nguyen, J. T., Barbisin, M., Xu, N. L., Mahuvakar, V. R., Andersen, M. R., Lao, K. Q., Livak, K. J., & Guegler, K. J. (2005). Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res*. 33, e179.
- Cissell, K. A., & Deo, S. K. (2009). Trends in microRNA detection. *Anal Bioanal Chem*. 394, 1109-1116.
- Dutta, R., Chomyk, A. M., Chang, A., Ribaudo, M. V., Deckard, S. A., Doud, M. K., Edberg, D. D., Bai, B., Li, M., Baranzini, S. E., Fox, R. J., Staugaitis, S. M., Macklin, W. B., & Trapp, B. D. (2013). Hippocampal demyelination and memory dysfunction are associated with increased levels of the neuronal microRNA miR-124 and reduced AMPA receptors. *Ann Neurol*. 73, 637-645.
- Fabian, M. R., & Sonenberg, N. (2012). The mechanics of miRNA-mediated gene silencing: a look under the hood of miRISC. *Nat Struct Mol Biol*. 19, 586-593.
- Gascon, E., Lynch, K., Ruan, H., Almeida, S., Verheyden, J. M., Seeley, W. W., Dickson, D. W., Petrucelli, L., Sun, D., Jiao, J., Zhou, H., Jakovcevski, M., Akbarian, S., Yao, W. D., & Gao, F. B. (2014). Alterations in microRNA-124 and AMPA receptors contribute to social behavioral deficits in frontotemporal dementia. *Nat Med*. 20, 1444-1451.
- Gentile, G., Morello, G., La Cognata, V., Guarnaccia, M., Conforti, F. L., & Cavallaro, S. (2022). Dysregulated miRNAs as Biomarkers and Therapeutical Targets in Neurodegenerative Diseases. *J Pers Med*. 12,
- Ghafouri-Fard, S., Khoshbakht, T., Hussen, B. M., Taheri, M., Ebrahimzadeh, K., & Noroozi, R. (2022). The emerging role of long non-coding RNAs, microRNAs, and an accelerated epigenetic age in Huntington's disease. *Front Aging Neurosci*. 14, 987174.
- Gilbert, J., Shu, S., Yang, X., Lu, Y., Zhu, L. Q., & Man, H. Y. (2016). β -Amyloid triggers aberrant over-scaling of homeostatic synaptic plasticity. *Acta Neuropathol Commun*. 4, 131.
- Grosjean, S., Pachón-Angona, I., Dawra, M., Refouvelet, B., & Ismaili, L. (2022). Multicomponent reactions as a privileged tool for multitarget-directed ligand strategies in Alzheimer's disease therapy. *Future Med Chem*. 14, 1583-1606.
- Hou, Q., Ruan, H., Gilbert, J., Wang, G., Ma, Q., Yao, W. D., & Man, H. Y. (2015). MicroRNA miR124 is required for the expression of homeostatic synaptic plasticity. *Nat Commun*. 6, 10045.
- Huntzinger, E., & Izaurralde, E. (2011). Gene silencing by microRNAs: contributions of translational repression and mRNA decay. *Nat Rev Genet*. 12, 99-110.

- Ibrahim, M. M., & Gabr, M. T. (2019). Multitarget therapeutic strategies for Alzheimer's disease. *Neural Regen Res.* 14, 437-440.
- Izaurralde, E. (2015). GENE REGULATION. Breakers and blockers—miRNAs at work. *Science.* 349, 380-382.
- Kato, Y., Sawata, S. Y., & Inoue, A. (2010). A lentiviral vector encoding two fluorescent proteins enables imaging of adenoviral infection via adenovirus-encoded miRNAs in single living cells. *J Biochem.* 147, 63-71.
- Kawakami, K., & Noda, T. (2004). Transposition of the Tol2 element, an Ac-like element from the Japanese medaka fish *Oryzias latipes*, in mouse embryonic stem cells. *Genetics.* 166, 895-899.
- Keaveney, M. K., Rahsepar, B., Tseng, H. A., Fernandez, F. R., Mount, R. A., Ta, T., White, J. A., Berg, J., & Han, X. (2020). CaMKII α -Positive Interneurons Identified via a microRNA-Based Viral Gene Targeting Strategy. *J Neurosci.* 40, 9576-9588.
- Keaveney, M. K., Tseng, H. A., Ta, T. L., Gritton, H. J., Man, H. Y., & Han, X. (2018). A MicroRNA-Based Gene-Targeting Tool for Virally Labeling Interneurons in the Rodent Cortex. *Cell Rep.* 24, 294-303.
- Kim, S. W., Li, Z., Moore, P. S., Monaghan, A. P., Chang, Y., Nichols, M., & John, B. (2010). A sensitive non-radioactive northern blot method to detect small RNAs. *Nucleic Acids Res.* 38, e98.
- Kosik, K. S., & Krichevsky, A. M. (2005). The Elegance of the MicroRNAs: A Neuronal Perspective. *Neuron.* 47, 779-782.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., & Tuschl, T. (2002). Identification of tissue-specific microRNAs from mouse. *Curr Biol.* 12, 735-739.
- Lee, C. Y., Ryu, I. S., Ryu, J. H., & Cho, H. J. (2021). miRNAs as Therapeutic Tools in Alzheimer's Disease. *Int J Mol Sci.* 22,
- Liu, N., Landreh, M., Cao, K., Abe, M., Hendriks, G. J., Kennerdell, J. R., Zhu, Y., Wang, L. S., & Bonini, N. M. (2012). The microRNA miR-34 modulates ageing and neurodegeneration in *Drosophila*. *Nature.* 482, 519-523.
- Macosko, E. Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A. R., Kamitaki, N., Martersteck, E. M., Trombetta, J. J., Weitz, D. A., Sanes, J. R., Shalek, A. K., Regev, A., & McCarroll, S. A. (2015). Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell.* 161, 1202-1214.
- Maiorano, N. A., & Mallamaci, A. (2010). The pro-differentiating role of miR-124: indicating the road to become a neuron. *RNA Biol.* 7, 528-533.
- Makeyev, E. V., Zhang, J., Carrasco, M. A., & Maniatis, T. (2007). The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell.* 27, 435-448.
- Malik, S. Z., Maronski, M. A., Dichter, M. A., & Watson, D. J. (2012). The use of specific AAV serotypes to stably transduce primary CNS neuron cultures. *Methods Mol Biol.* 846, 305-319.
- Millan, M. J. (2017). Linking deregulation of non-coding RNA to the core pathophysiology of Alzheimer's disease: An integrative review. *Prog Neurobiol.* 156, 1-68.
- Namkung, H., Yukitake, H., Fukudome, D., Lee, B. J., Tian, M., Ursini, G., Saito, A., Lam, S., Kannan, S., Srivastava, R., Niwa, M., Sharma, K., Zandi, P., Jaaro-

- Peled, H., Ishizuka, K., Chatterjee, N., Haganir, R. L., & Sawa, A. (2023). The miR-124-AMPA pathway connects polygenic risks with behavioral changes shared between schizophrenia and bipolar disorder. *Neuron*. 111, 220-235.e9.
- Nguyen, T. P. N., Kumar, M., Fedele, E., Bonanno, G., & Bonifacino, T. (2022). MicroRNA Alteration, Application as Biomarkers, and Therapeutic Approaches in Neurodegenerative Diseases. *Int J Mol Sci*. 23,
- Podnar, J., Deiderick, H., Huerta, G., & Hunicke-Smith, S. (2014). Next-Generation Sequencing RNA-Seq Library Construction. *Curr Protoc Mol Biol*. 106, 4.21.1-19.
- Rosenberg, A. B., Roco, C. M., Muscat, R. A., Kuchina, A., Sample, P., Yao, Z., Graybuck, L. T., Peeler, D. J., Mukherjee, S., Chen, W., Pun, S. H., Sellers, D. L., Tasic, B., & Seelig, G. (2018). Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding. *Science*. 360, 176-182.
- Salta, E., & De Strooper, B. (2017). microRNA-132: a key noncoding RNA operating in the cellular phase of Alzheimer's disease. *FASEB J*. 31, 424-433.
- Sano, M., Ohtaka, M., Iijima, M., Nakasu, A., Kato, Y., & Nakanishi, M. (2017). Sensitive and long-term monitoring of intracellular microRNAs using a non-integrating cytoplasmic RNA vector. *Sci Rep*. 7, 12673.
- Seyedaghamiri, F., Rajabi, M., & Mohaddes, G. (2023). Targeting Novel microRNAs in Developing Novel Alzheimer's Disease Treatments. *Neurochem Res*. 48, 26-38.
- Song, Y., Xu, Z., & Wang, F. (2020). Genetically Encoded Reporter Genes for MicroRNA Imaging in Living Cells and Animals. *Mol Ther Nucleic Acids*. 21, 555-567.
- Su, Q., Sena-Estevés, M., & Gao, G. (2020). Production of Recombinant Adeno-Associated Viruses (rAAVs) by Transient Transfection. *Cold Spring Harb Protoc*. 2020, 095596.
- Tran, K. K. N., Wong, V. H. Y., Hoang, A., Finkelstein, D. I., Bui, B. V., & Nguyen, C. T. O. (2023). Retinal alpha-synuclein accumulation correlates with retinal dysfunction and structural thinning in the A53T mouse model of Parkinson's disease. *Front Neurosci*. 17, 1146979.
- Urbanek, M. O., Nawrocka, A. U., & Krzyzosiak, W. J. (2015). Small RNA Detection by in Situ Hybridization Methods. *Int J Mol Sci*. 16, 13259-13286.
- Valencia-Sanchez, M. A., Liu, J., Hannon, G. J., & Parker, R. (2006). Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev*. 20, 515-524.
- Visvanathan, J., Lee, S., Lee, B., Lee, J. W., & Lee, S. K. (2007). The microRNA miR-124 antagonizes the anti-neural REST/SCP1 pathway during embryonic CNS development. *Genes Dev*. 21, 744-749.
- Wang, X., Liu, D., Huang, H. Z., Wang, Z. H., Hou, T. Y., Yang, X., Pang, P., Wei, N., Zhou, Y. F., Dupras, M. J., Calon, F., Wang, Y. T., Man, H. Y., Chen, J. G., Wang, J. Z., Hébert, S. S., Lu, Y., & Zhu, L. Q. (2018). A Novel MicroRNA-124/PTPN1 Signal Pathway Mediates Synaptic and Memory Deficits in Alzheimer's Disease. *Biol Psychiatry*. 83, 395-405.
- Wang, X. W., Hu, L. F., Hao, J., Liao, L. Q., Chiu, Y. T., Shi, M., & Wang, Y. (2019). A microRNA-inducible CRISPR-Cas9 platform serves as a microRNA sensor and cell-type-specific genome regulation tool. *Nat Cell Biol*. 21, 522-530.

- Xue, Q., Yu, C., Wang, Y., Liu, L., Zhang, K., Fang, C., Liu, F., Bian, G., Song, B., Yang, A., Ju, G., & Wang, J. (2016). miR-9 and miR-124 synergistically affect regulation of dendritic branching via the AKT/GSK3 β pathway by targeting Rap2a. *Sci Rep.* 6, 26781.
- Yao, L., Zhu, Z., Wu, J., Zhang, Y., Zhang, H., Sun, X., Qian, C., Wang, B., Xie, L., Zhang, S., & Lu, G. (2019). MicroRNA-124 regulates the expression of p62/p38 and promotes autophagy in the inflammatory pathogenesis of Parkinson's disease. *FASEB J.* 33, 8648-8665.
- Yekta, S., Shih, I. H., & Bartel, D. P. (2004). MicroRNA-directed cleavage of HOXB8 mRNA. *Science.* 304, 594-596.
- Zhao, Y., Zheng, Q., Hong, Y., Gao, Y., Hu, J., Lang, M., Zhang, H., Zhou, Y., Luo, H., Zhang, X., Sun, H., Yan, X. X., Huang, T. Y., Wang, Y. J., Xu, H., Liu, C., & Wang, X. (2023). β_2 -Microglobulin coaggregates with A β and contributes to amyloid pathology and cognitive deficits in Alzheimer's disease model mice. *Nat Neurosci.*
- Zhou, L., Li, X., Liu, Q., Zhao, F., & Wu, J. (2011). Small RNA transcriptome investigation based on next-generation sequencing technology. *J Genet Genomics.* 38, 505-513.