

Predominant heterotrophic diazotrophic bacteria are involved in Sargassum proliferation in the great Atlantic Sargassum Belt

M Léger-Pigout, E Navarro, F Menard, S Ruitton, F Le Loc'h, S Guasco, J-M Munaron, D Thibault, T Changeux, S Connan, et al.

► To cite this version:

M Léger-Pigout, E Navarro, F Menard, S Ruitton, F Le Loc'h, et al.
. Predominant heterotrophic diazotrophic bacteria are involved in Sargas
sum proliferation in the great Atlantic Sargas
sum Belt. The International Society of Microbiologial Ecology Journal, 2024, 10.1093/ismejo/wrad026 . hal-04421726

HAL Id: hal-04421726 https://amu.hal.science/hal-04421726

Submitted on 6 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



https://doi.org/10.1093/ismejo/wrad026 Advance access publication: 10 January 2024 Original Article

Predominant heterotrophic diazotrophic bacteria are involved in Sargassum proliferation in the Great Atlantic Sargassum Belt

Matéo Léger-Pigout¹, Elisabeth Navarro¹, Frédéric Ménard¹, Sandrine Ruitton¹, François Le Loc'h², Sophie Guasco¹,

Jean-Marie Munaron², Delphine Thibault¹, Thomas Changeux¹, Solène Connan², Valérie Stiger-Pouvreau², Thierry Thibaut¹,

Valérie Michotey^{1,*}

¹Aix Marseille Univ, Université de Toulon, CNRS, IRD, MIO, Marseille, France ²Univ Brest, CNRS, IRD, Ifremer, LEMAR, F-29280 Plouzane, France

*Corresponding author: Valérie Michotey, MIO, Bat 26, Campus de Luminy, Case 901, Av de Luminy, 13288 Marseille cedex 09, France. Email: valerie.michotey@univ-amu.fr

Abstract

Since 2011, the Caribbean coasts have been subject to episodic influxes of floating *Sargassum* seaweed of unprecedented magnitude originating from a new area "the Great Atlantic *Sargassum* Belt" (GASB), leading in episodic influxes and mass strandings of floating *Sargassum*. For the biofilm of both holopelagic and benthic *Sargassum* as well as in the surrounding waters, we characterized the main functional groups involved in the microbial nitrogen cycle. The abundance of genes representing nitrogen fixation (*nifH*), nitrification (*amoA*), and denitrification (*nosZ*) showed the predominance of diazotrophs, particularly within the GASB and the Sargasso Sea. In both location, the biofilm associated with holopelagic *Sargassum* harboured a more abundant proportion of diazotrophs than the surrounding water. The mean δ^{15} N value of the GASB seaweed was very negative ($-2.04\%_0$), and lower than previously reported, reinforcing the hypothesis that the source of nitrogen comes from the nitrogen-fixing activity of diazotrophs within this new area of proliferation. Analysis of the diversity of diazotrophic communities revealed for the first time the predominance of heterotrophic diazotrophic bacteria belonging to the phylum *Proteobacteria* in holopelagic *Sargassum* biofilms. The *nifH* sequences belonging to Vibrio genus (*Gammaproteobacteria*) and Filomicrobium sp. (Alphaproteobacteria) were the most abundant and reached, respectively, up to 46.0% and 33.2% of the community. We highlighted the atmospheric origin of the nitrogen used during the growth of holopelagic *Sargassum* within the GASB and a contribution of heterotrophic nitrogen-fixing bacteria to a part of the *Sargassum* proliferation.

Keywords: Sargassum, N2 fixation, nitrification, denitrification, GASB, isotopy, Vibrio spp

Introduction

Since 2011, blooms of the holopelagic brown macroalgae Sargassum spp. have been observed in the tropical North Atlantic Ocean, causing significant beach strandings along the Caribbean coasts. The presence of Sargassum is a permanent feature of the North Atlantic, where massive quantities are trapped in the Sargasso Sea by large-scale hydrodynamics. Holopelagic Sargassum has flourished naturally in this area despite the nutrient-poor and low-productivity characteristics of the surrounding waters [1]. Several factors could explain this apparent paradox: first, Sargassum growth is driven by new production in the neritic waters of the western North Atlantic Ocean and the Gulf of Mexico, where mutualistic relationships with fish could contribute to nutrient supply [2]. Second, seasonal transport of nutrient-rich and productive Sargassum from the Gulf of Mexico is facilitated by largescale currents such as the Loop Current and the Gulf Stream, which enter the Sargasso Sea [3]. Third, nitrogen fixation by the Cyanobacteria Dichothrix fucicola, an epiphyte on Sargassum, has been shown to provide a significant portion (2%-32%) of the nitrogen demand for the holopelagic Sargassum in the Sargasso Sea

[4, 5]. Recently N fixation rates associated with Sargassum ranging from 0 to 30916 μ mol.N.m⁻².d⁻¹ have been recorded in the North Atlantic [6]. This nitrogen fixation contributes to variable δ^{15} N enrichment in Sargassum tissues, ranging from approximately 8‰ to -2% [7, 8].

Sargassum stranding occurs along the Gulf of Mexico coast, resulting in negative economic [9-13] and environmental [13-15] impacts. The development of Sargassum in this region has been attributed to increasing nitrogen inputs from various land-based sources, including the Mississippi River and its tributaries [3, 16]. The higher δ^{15} N enrichment of Sargassum in the Gulf of Mexico (in the range of 5‰–8‰) compared to that observed in the Sargasso Sea may be related to the urbanization of the Texas coast and the Mississippi River plume [8].

In 2010–2011, a significant biomass of *Sargassum* was discovered in a new region of the tropical Atlantic Ocean south of the Sargasso Sea, where it had never previously occurred in such abundance [17, 18]. Since then, the seaweed has flourished, forming a new consolidated region called the "Great Atlantic *Sargassum* Belt" (GASB), which extends from the coasts of Africa to South and Central America. These drifting seaweeds are carried by

Received 20 September 2023. Revised: 27 November 2023. Accepted: 8 December 2023

© The Author(s) 2024. Published by Oxford University Press on behalf of the International Society for Microbial Ecology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

ocean currents and winds, aggregate at the surface to form visible rafts that can be detected by satellite remote sensing across the transatlantic region. As a result, large quantities, reaching up to 20 million tons, regularly wash up on the coasts of the Caribbean islands, northern Brazil, Guyana, and West Africa [19]. This represents an emerging threat to areas already under anthropogenic pressure. As in the Gulf of Mexico, the rapid decomposition of stranded *Sargassum* in these regions poses significant health risks to local populations [20] and economy [21] due to the release of hydrogen sulphide (H₂S), a toxic and corrosive gas [22].

These multiregional transboundary *Sargassum* influxes in the GASB do not appear to result from a single or simple cause, but rather from complex combination of causal factors. These include higher water temperatures [23, 24], nutrient inputs from rivers in the region, such as the Amazon [8], and upwelling of nutrient-rich deep water [19, 25]. Nevertheless, these hypotheses do not fully explain the extent of proliferation [23, 26-29], and significant gaps remain in our knowledge and understanding. For example, the potential role of *Sargassum* microbial biofilms in promoting algal growth has not been thoroughly investigated in the GASB.

In a 2020 study [30], an analysis of samples from the GASB and revealed the presence of four distinct *Sargassum*-associated microbial biofilms, with only one shared with the Gulf of Mexico counterpart [30, 31]. A subsequent study in 2023 [32] also showed biogeographic patterning of the holopelagic *Sargassum* microbiome and identified biomarker genera indicative of *Sargassum natans* I/VIII and *Sargassum* fluitans III. This suggests that the new location may select for different microbial members within the *Sargassum* biofilm. As nitrogen was identified as a potential limiting factor for *Sargassum* growth, we thus investigate the nitrogen cycle and the interactions between *Sargassum* and its prokaryotic biofilm partners in the GASB.

As a result, our study quantified the contribution of key microbial members of the nitrogen cycle, such as diazotrophs, nitrifiers, and denitrifiers, responsible for the entry, transformation, and exit of assimilable nitrogen, from holopelagic *Sargassum* and its surrounding waters. This large-scale study extended from the GASB to the Caribbean Islands and the southern Sargasso Sea, including samples of benthic *Sargassum* in the Caribbean. The study focused on the diversity of diazotrophs and analysed their relationships with the δ^{15} N values of the seaweed and the particulate organic matter (POM), including phytoplankton, in the adjacent water.

This approach aimed to investigate the relative contributions of these microbial members to the growth of *Sargassum* in the GASB and Caribbean coastal zones and ultimately identified additional key microbial players that could contribute to the proliferation of *Sargassum*.

Materials and methods Sampling campaigns

Twenty sampling stations were set up during two sea cruises and one coastal mission in the French West Indies (Fig. 1). The Western Atlantic-Sargassum expedition (https://doi.org/10.17600/17004300) took place aboard the N/O ANTEA from 19 June to 13 July 2017 and explored the GASB as well as the Sargasso Sea (25°N). The Transatlantic-Sargassum expedition (https://doi. org/10.17600/17016900) took place aboard the M/V YERSIN from 6 to 24 October 2017 from the Cape Verde Islands, crossing the Atlantic between 8 and 12°N to the island of Martinique within the GASB. The SAVE-C coastal sampling period along the coasts of Martinique and Guadeloupe took place from 7 to 17 July 2021. Despite the fact that sampling missions were performed at different years and seasons (Antea and Yersin in 2017 and SAVE-C in 2021, and spring/summer for Antea and SAVE-C and Autumn for Yersin), water temperatures were around 28°C and showed less than 1°C variation (Antea 27.6°C to 28.7°C, Yersin 27.4°C to 27.7°C, and Save-C 27.42°C to 28.28°C). At each station, apical sections of stems with leaves and small bladders (pneumatocysts) of the three holopelagic morphotypes (*S. natans* I and VIII and *S. fluitans* III identified as described previously [33]) as well as water samples surrounding the rafts were collected in at least triplicate. The typology of the sampled *Sargassum* rafts [34] was also recorded.

To analyse the biofilm and associated microbial communities growing on Sargassum (n = 159), approximately 5 g wet weight of Sargassum was placed in a 100 -cm³ cup, filled with 70 cm³ of sterile seawater and sonicated for 30 s (46 kHz, 30 W) to dissociate the bacterial biofilm from its support [30]. The water containing the dissociated biofilm was then filtered through 0.22 - μ m filters (nitrocellulose membrane, 47 mm), and the filters were stored at -20° C. For the microbial communities in the surrounding waters, 4 l of surface water was collected adjacent to the sampled Sargassum using a clean bucket. Planktonic microorganisms were recovered by filtering the water through a 0.22 - μ m nitrocellulose membrane (47 mm, n = 83) in at least in triplicate for each station. The filters were stored at -20° C until further analysis.

Nitrogen stable isotope analysis (δ^{15} N) was conducted on *Sargassum* thallus from the three morphotypes with at least three samples per morphotype. The samples (n=262) were promptly stored in silica gel. At a particular location, stems, leaves, and bladders were examined separately. During the sampling period in Martinique and Guadeloupe, we tested the impact of biofilm microbial communities on the isotopic composition by analyzing samples with (n=45) and without (n=34) biofilm, which has been removed through sonication. The isotopic composition of the POM, which includes planktonic microorganisms, was acquired by filtering at least 20 l of the surrounding water in triplicate through preweighed Whatman GF/F glass microfiber 0.7- μ m filters (precombusted at 500°C for 4 h, n = 110). The filters were subsequently dried and stored in silica gel until further analysis.

Microbial analysis

To better characterize the nitrogen cycle in the Sargassum ecosystem of GASB compared to other areas, microbial analysis was performed on the frozen samples. The DNA collected on nitrocellulose membranes from microorganisms of the Sargassum biofilm as well as the water surrounding the Sargassum were extracted using a published protocol [30] for the Antea and Yersin cruises samples, and DNAeasy®PowerLyser®PowerSoil®Kit (Qiagen) was used for SAVE-C cruise samples. Diazotrophs, nitrifiers, and denitrifiers were then studied through the relative abundance of a selection of their marker genes (nifH, amoA, and nosZ, respectively) relatively to ribosomal gene (16S rRNA genes). qPCR were performed on DNA from all biofilms and surrounding water samples. Denitrifiers quantification (nosZ gene) was performed with the primers nosZ 2F [35]—nosZ 1897R [36]. Bacterial nitrifiers quantification (amoA gene) was performed with the primers amoA-1F—amoA-2R [37] and the primers arc-amoA F100 mod-arc-amoA R336 mod [38] were used to amplify the archeal *amoA* gene. The mixes (20 μ l) used in these qPCRs were composed of EvaGreen 2X (10 μ l Bio-Rad[®]), primer F/R (0.5 μ l), DNA (2 μ l), and H₂O (7 μ l). We used the following program during qPCR of the genes nosZ, bacterial amoA, and archeal amoA: an initial denaturation (98°C for 5 min, 2 min, 2 min, respectively) followed by 40 cycles each comprising



Figure 1. Location of sampling stations (Sargassum and surrounding waters). ANTEA (June 19 to July 13, 2017, S-stations), YERSIN (10–24 October 2017, Y-stations), SAVE-C (7–17 July 2021, M- or G-stations) campaigns. Holopelagic Sargassum samples are represented by dots and benthic samples by triangles.

a denaturation step (98°C for 15 s-5 s, respectively), hybridization (64°C-58°C-62°C for 15 s, respectively), elongation (72°C for 40 s-1 min-1 min, respectively), and a final elongation (95°C for 10 s). For qPCR of the nifH gene, the primers used commonly for the nested PCRs used in the *nifH* gene amplicon sequencing analysis [39] were not compatible with this analysis. Instead, we used the PolF/PolR primers [40], frequently used in marine and terrestrial environments. Validation of PolF/PolR primers coverage against GenBank nr database was performed using all combinations offered by the degenerate sequences of these two primers. A coverage of all the diazotroph taxa was observed. Moreover, these primers covered more than 90% of the relative abundance obtained after nifH gene amplicon sequencing analysis (Supplementary Table 1). We also tested their amplification efficiency on proteobacterial and cyanobacterial fragments obtained in our study and that had been previously inserted into plasmids. In these tests, the calibration curve exhibited good characteristics $(R^2 = 0.99; E = 91.8\%)$ and was used as the standard range for the entire analysis (Supplementary Fig. 1). The mixes (20 μ l) used in the nifH-qPCR were composed of EvaGreen 2X (10 μ l, Bio-Rad[®]), PolF/PolR (0.5 μ l), DNA (2 μ l), and H₂O (7 μ l). We used the following program during qPCR: an initial denaturation (98°C for 2 min) followed by 45 cycles each comprising a denaturation step (98°C for 5 s), hybridization (55°C for 15 s), elongation (72°C for 15 s), and finally a melting curve (from 65°C to 95°C with an increment of 0.5°C every 5 s). Each qPCR was validated following the observation of a correct melting curve, considering the large variability in nifH gene size [41]. Quantification of the 16S rRNA genes was performed as described previously [30].

The diversity of diazotroph was studied at each station (biofilm and surrounding water n=35) after pooling replicate DNA extracts. For nifH gene amplicon sequence analysis, MiSeq (Illumina) 2×250 bp sequencing of the nifH gene was performed (Genewiz®) following nested PCR amplification, with nifH1/nifH2 and nifH3/nifH4 sets [39] and sequencing. Primer deletion, quality filtering, dereplication, error correction, sample inference,

merging of paired reads, chimaera removal, and taxonomic classification of raw sequences were performed using R software (4.1.2) and dada2 (1.22.0) pipeline [42]. Approximately 5000000 reads were obtained from the 35 samples (145000 reads per sample on average). A first taxonomic assignment was performed against a nifH database (June 2017 version) [43] in dada2 format (https://doi.org/10.5281/zenodo.3958370). As many amplicon sequence variants (ASVs) were not assigned or did not have a precise taxonomic assignment for this study, we enriched the database with assigned sequences specific to our dataset. Each ASV that did not have a taxonomic assignment at the class level was subjected to BLAST analysis against the GenBank nr database. Sequences that matched with ASVs at a high percentage identity (>85%) were integrated into the initial database. A second taxonomic analysis was performed with this in-house database. In total, 942 ASVs were identified via this analysis (Supplementary Data1). However, this method was not able to assign all the ASVs to the class level. We therefore performed the class phylogeny of all unassigned ASVs representing at least 1% of each sample using Clustal Omega alignment (https:// www.ebi.ac.uk/Tools/msa/clustalo/) and NJ tree constructed using SeaWiew software (https://doua.prabi.fr/software/seaview) (Supplementary Fig. 2). Raw sequence data are available at the NCBI Sequence Read Archive under bio project accession #PRJNA1017983.

Determination of δ^{15} N in Sargassum and POM samples

For pieces of *Sargassum*, approximately 2 g of sample dried powder was weighed in tin capsules for isotopic analysis. For POM, samples were duplicated: Only the subsamples that were not treated were analysed for δ^{15} N. Stable isotope analyses were performed by continuous flow on a Flash EA 2000 elemental analyser (Thermo Scientific, Milan, Italy) coupled to a Delta V Plus, isotope ratio mass spectrometer (Thermo Scientific, Bremen, Germany) at the Pôle Spectrométrie Océan (IUEM-UBO, Plouzané, France). The reference materials (USGS-61, USGS-62, and USGS-63) and an in house standard Thermo-Acetanilide were used for isotopic corrections and to assign the data to the appropriate isotopic scale: USGS-61 (certified values: $\delta^{15}N = -2.87\% \pm 0.04\%$ and measured values: $\delta^{15}N = -2.89\% \pm 0.06\%$), USGS-62 (certified values: $\delta^{15}N = +20.17\% \pm 0.06\%$ and measured values: $\delta^{15}N = +20.17\% \pm 0.08\%$), and USGS-63 (certified values: $\delta^{15}N = +37.83\% \pm 0.06\%$ and measured values: $\delta^{15}N = +37.83\% \pm 0.06\%$ and measured values: $\delta^{15}N = +37.83\% \pm 0.06\%$ and measured values: $\delta^{15}N = +37.84\% \pm 0.10\%$). Results were reported in the δ unit notation and expressed as parts per thousand relative to the international standards atmospheric N₂ for nitrogen. Analytical precision based on replicate measurements (after every five samples) of Thermo-Acetanilide was <0.1 for $\delta^{15}N$ values.

Statistics

All statistics were performed using R software (4.1.3). To define how the abundances of the nitrogen cycle members and the $\delta^{15}N$ varied between the different microbiomes, the normality of the groups was first tested (stats package, 4.1.2) using Shapiro-Wilk tests. Then nonparametric Kruskal-Wallis analysis followed by Nemenyi tests (PMCMRplus package, 1.9.6), Welch's t-tests, and Wilcoxon tests were performed. Plots and associated 95% confidence intervals of the mean (2000 bootstraps) were drawn using the *qqplot2* package (3.4.2). To assess the variation of diazotrophic communities according to microbiomes, we first replaced the zeros in the ASV table using the zCompositions package. The ASV table was then transformed into a compositional log-ratio (clr) matrix to mitigate the compositional nature of the data. An Aitchison distance matrix was then computed using the phyloseq package (1.38.0). The NMDS analysis was performed on the Aitchison distance matrix using the vegan package (2.6.4), employing a configuration with two dimensions (k = 2) and a maximum of 500 iterations (trymax = 500), with an initial configuration generated after 200 tries (try=200). In addition to NMDS, a Hierarchical Ascendant Classification (HAC) based on the Aitchison distance matrix was performed (method = complete) using the stats package (4.1.2). The dendrograms obtained were cut into four clusters using the cutree function in the stats package (4.1.2). Dunn's index was calculated using the package fpc (2.2.10).

Results

Relative abundance of bacteria involved in nitrogen cycle processes

In the holopelagic *Sargassum* biofilm from the offshore samples (GASB and Sargasso Sea), diazotrophs (nifH/16S rRNA genes ratio) were found to comprise 2% \pm 1% and were significantly dominant over denitrifiers and nitrifiers (approximately 10³ to 10⁵ times higher, P < .001) (Fig. 2A). However, biofilm composition underwent significant changes in coastal samples, and varying trends were observed among the microbial communities. Diazotrophs exhibited a significant 10-fold decrease (P < .05) in abundance between offshore and coastal samples, reaching approximatively 0.1%. In contrast, nitrifiers and denitrifiers showed a substantial increase, respectively, 10^2 - and 10^3 -fold higher (P < .001). Consequently, in the holopelagic Sargassum biofilm collected from the coastal region of Martinique and Guadeloupe, denitrifiers became prevalent over diazotrophs (P < .001) and reached $9.7\% \pm 4.1\%$ (nosZ/16S rRNA genes ratio). Within samples from the coastal area, the relative abundance of diazotrophs, nitrifiers, and denitrifiers in the holopelagic Sargassum biofilm was higher compared to that in the biofilm of benthic Sargassum collected on the same island (Fig. 2B-D). To investigate possible connections

between holopelagic Sargassum and its biofilm, a comparison between relative abundance of the analysed genes collected from Sargassum biofilm and their surrounding water was performed. In offshore samples, relative abundance of diazotrophs or nitrifiers community between Sargassum biofilm and surrounding water showed significant differences (P < .05; Fig. 2B–D) in contrast to denitrifiers (Fig. 2C). In coastal areas, no difference was observed between the relative abundance of the analysed genes retrieved from holopelagic Sargassum or its surrounding waters. Consequently, conditions encountered within biofilms of holopelagic Sargassum in offshore samples (GASB and Sargasso Sea) seem to favour diazotroph and nitrifier development compared to the surrounding waters. This positive effect seems to disappear in coastal areas. To determine whether raft size, Sargassum morphotype, or geographic location affect biofilm composition, the relative abundances of the selected genes were studied among our extensive set of biofilm samples (Supplementary Data 2). Our results indicate that neither the morphotypes of holopelagic Sargassum nor raft types had a significant impact on the relative abundance of nifH, nosZ, and amoA genes in the biofilm within the GASB (Supplementary Fig. 3, P > .05). Gene abundance was found to be similar (P > .05; Fig. 2) in offshore samples.

Determination of the source of nitrogen in Sargassum and POM of their surrounding waters

Isotopic analysis of δ ¹⁵N was conducted to examine the contribution of nitrogen by diazotrophs (Supplementary Data 3). Measurements of δ^{15} N were performed on Sargassum and POM (including phytoplankton) from the same sample set (holopelagic Sargassum from offshore and coastal area, benthic Sargassum, n = 262). Analyses were performed on Sargassum with their entire biofilm or mostly removed by sonication, and on different parts of the seaweed at sampling stations located in the centre of the GASB (leave, stem, and bladder). The δ^{15} N value of offshore holopelagic Sargassum (with biofilm) was extremely low (-2.04‰ on average) with no differences observed between GASB and Sargasso Sea samples (Fig. 3A). In the coastal zone, $\delta^{15}N$ of holopelagic Sargassum (with biofilm) increased to reach about -1‰. Sonication of holopelagic Sargassum, to disassociate the biofilm, did not have any significant effect on $\delta^{15}N$ of the seaweed (P > .05). Furthermore, in the centre of the GASB (Y11 station), δ^{15} N of the stem, leaves, and bladders of the holopelagic Sargassum presented neither significant difference between them nor with the entire seaweed (P > .05) reflecting a common nitrogen source (Fig. 3B). Overall δ^{15} N of holopelagic Sargassum (-2.04‰ on average) are significantly lower than those measured for the corresponding POM (5.06% on average) or that of benthic Sargassum (δ^{15} N of about 0‰) (Fig. 3A). These results suggest that fixed N₂ by microbial biofilm accounts for an important proportion of the assimilated nitrogen of the holopelagic Sargassum of GASB, with a constant source along the growth of the different parts of the thallus. However, the amount of fixed nitrogen decreased upon reaching the coastal Caribbean region.

Biofilm containing an original diazotrophic community structure

Analyses of *nifH* gene amplicon sequencing revealed differences in diazotroph compositions between holopelagic and benthic *Sargassum* biofilm (BS-B), or corresponding surrounding waters (Figs 4 and 5; Supplementary Fig. 3). In the offshore, holopelagic *Sargassum* biofilm communities were predominantly composed of noncyanobacterial diazotrophs (NCDs, $76.1\% \pm 13.2\%$), with *Cyanobacteria* representing only $23.9\% \pm 13.2\%$ of the diazotroph



Figure 2. Log₁₀ evolution of the ratio of *nifH*, *nosZ*, and *amoA* gene abundances to 16S rRNA genes abundance in different microbiomes. For each graph, samples are categorized according to their origins: biofilm or water, pelagic or benthic, open ocean or coastal, samples. Each grey point corresponds to one data item. The 95% confidence intervals of the mean are shown. (A) Relative abundance of the *nifH*, *nosZ*, and *amoA* genes in the *Sargassum* biofilm. (B) Relative abundance of the *nifH* gene across microbiomes. (C) Relative abundance of the *nosZ* gene across microbiomes. (D) Relative abundance of the *amoA* gene across microbiomes. SS, Sargasso Sea; MAR, Martinique; GUA, Guadeloupe.



Figure 3. Evolution of $\delta^{15}N(\%_0)$ as a function of the different sampled microbiomes. (A) Evolution of $\delta^{15}N(\%_0)$ as a function of different sample types: holopelagic and benthic *Sargassum*, and POM and between open ocean and coastal areas samples. (B) Evolution of $\delta^{15}N(\%_0)$ as a function of the different parts of the holopelagic *Sargassum* (all parts combined, bladders, leave, stem). GASB corresponds to the Great Atlantic *Sargassum* Belt. SS corresponds to Sargasso Sea. Each point corresponds to one data. The 95% confidence intervals of the mean are shown.

community. NCDs are dominated by Proteobacteria that reached on average $68.3\% \pm 19.3\%$ of the diazotroph community (Fig. 5; Supplementary Fig. 3). Additionally, there is a small percentage of Planctomycetes (4%) and Firmicutes phylum (1%). Among the NCDs, the two most abundant taxa (ASV) corresponded to a *Gammaproteobacteria* for which the closest sequence corresponds to Vibrio sp. (abundance among nifH up to 46.0%) and an Alphaproteobacteria that cluster with Filomicrobium sp. (abundance up to 33.2%). Among the cyanobacterial diazotroph of the holopelagic *Sargassum* biofilm, the two most taxa were affiliated to the genera Mastigocoleus $(1.2\% \pm 0.9\%$ in average, order Nostocales) and Hyella $(1.3\% \pm 1.4\%$ in average, order Pleurocapsales). The communities within the surrounding waters did not show such a marked difference between NCDs (essentially Proteobacteria) and cyanobacterial diazotrophs, they represented, respectively, $55.6\% \pm 16.6\%$ and $39.4\% \pm 16.1\%$. The two most abundant NCD's taxa belonged, for the first one to a nonidentified Proteobacteria $(10.5\% \pm 15.4\%)$ and for the second one to an Alcaligenes sp. $(8.5\% \pm 7.2\%)$. The two most abundant Cyanobacteria taxa in water were both affiliated to Trichodesmium sp. $(27.3\% \pm 15.6\%)$



Figure 4. NMDS and HAC analysis of relative abundance of diazotrophs per station according to microbiomes. Each point represents a station. The groups correspond to holopelagic *Sargassum* biofilm (HS-B) and associated water (HS-W), benthic *Sargassum* biofilm (BS-B) and associated water (BS-W). The NMDS analysis (S=0.125) was performed using an Aitchison distance matrix. The four clusters obtained with a HAC (Dunn index = 0.59; Supplementary Fig. 5) are represented by the coloured envelopes.



Figure 5. Relative abundances (%) of diazotroph classes in relation to microbiomes. The microbiomes shown correspond to the holopelagic *Sargassum* biofilm (HS-B) and associated water (HS-W), and the benthic *Sargassum* biofilm (BS-B) and associated water (BS-W). NI, not identified.

and $9.6\% \pm 11.4\%$, respectively). Dichothrix fucicola populations appeared as minor member of the diazotroph community, and their contribution ranged between 1% and 6.5% (S8 and S10 stations, Supplementary Fig. 4) in the GASB. The biofilm of benthic Sargassum showed a diazotroph community mainly dominated by Proteobacteria, representing up to 75% in the biofilm of Sargassum (Supplementary Fig. 3).

Discussion

Because Sargassum could be found in oligotrophic environments with low nitrogen concentrations [8], it is crucial to determine the various nitrogen sources utilized by these macroalgae and the underlying mechanisms. Our study reveals the significance of N_2 fixation compared to nitrification and denitrification in GASB, and this pattern changes when the holopelagic Sargassum arrives in neritic waters.

Previous research has identified N2 fixation activity associated with Sargassum from the Sargasso Sea and the Gulf of Mexico, by acetylene reduction assays [4-6, 44, 45] and N₂ assimilation by δ^{15} N analysis [8, 46, 47]. Our study demonstrates that in the entire GASB and in the southern region of the Sargasso Sea, holopelagic Sargassum spp. exhibit negative δ^{15} N values that are significantly lower than those measured for POM collected in their surrounding water. Our findings suggest a nitrogen source from fixed nitrogen by biofilm and exclude the involvement of free-living diazotrophs. Additionally, our study reveals the N source coming from diazotrophic nitrogen fixation activity of the biofilm in the new proliferation area. However, the δ^{15} N values are lower than those measured in previous studies conducted in the Sargasso Sea and the Gulf of Mexico [8, 46, 47] indicating a higher level nitrogen fixation activity and an increased involvement of diazotrophs in suppling nitrogen to Sargassum in GASB. Similar isotopic values $(\delta^{15}N)$ reported throughout the different parts of the Sargassum indicate a uniform nitrogen source during seaweed growth along its transatlantic journey. Nitrogen transfer efficiency between biofilm and Sargassum was demonstrated by the identical $\delta^{15}N$ values of Sargassum both with and without its biofilm.

Despite their ecological importance, the diversity of the diazotroph community associated with Sargassum has been poorly characterized until now. In microscopic studies from the 1980s, Cyanobacteria including Dichothrix fucicola and others belonging to the genus Calothrix were identified [4]. Since then, Cyanobacteria have been considered the main diazotrophs and the primary contributors to nitrogen availability for Sargassum [4, 5]. Our findings from the GASB demonstrate that this idea is overly restrictive. Heterotrophic Proteobacteria, not Cyanobacteria, were the predominant diazotrophs in Sargassum biofilms in the GASB. The previous phylogenetic studies of the holopelagic Sargassum biofilm (based on 16S rRNA genes), collected in the same area [30-32], and from beach stranding [48, 49] have also observed a low occurrence of Cyanobacteria. Additionally, this study demonstrates, for the first time, the prevalence of heterotrophic bacteria (NCD) in the diazotroph community of Sargassum biofilm. Diazotrophs played a major role in the nitrogen cycle of Sargassum, particularly in GASB and the Sargasso Sea, which are known for stronger oligotrophic conditions as compared to coastal areas, as the isotopic results have demonstrated. In the latter regions, diazotrophy reduced, and the abundance of denitrifiers increase, resulting in an agreement with previous findings regarding the increase in δ^{15} N [50].

Our results also demonstrate that the diversity of diazotrophic bacterial communities in offshore is distinct between *Sargassum* biofilms and surrounding waters, as previously reported [30, 31] based on 16S rRNA genes bacterial communities. The greater abundance of diazotrophs and higher prevalence of NCD in the holopelagic *Sargassum* biofilm compared to the surrounding waters in offshore suggest potential reciprocal benefits. Independent of the level of bacterial community analysis, in the open ocean, *Sargassum* biofilms are colonized by specific populations reinforcing the potential occurrence of close metabolite exchange links between seaweeds and their biofilms.

The source of carbon required for the metabolism of heterotrophic nitrogen fixing bacteria identified here remains unknown. The photo(hetero)trophs and organic matter degraders mainly composed the microbial community of holopelagic *Sargassum* [32] and could be one of carbon source. Furthermore, several studies have revealed that *Sargassum* spp. is a significant source of dissolved organic carbon [51, 52], could be directly metabolized by heterotrophic diazotrophs within their biofilm, and should be an example of mutualism relationships. Within the NCDs, an ASV clustering with Vibrio spp. *nifH* sequences was found to be among the most abundant, suggesting a biogeochemical role for this taxon in addition to a potential health risks in the area [30, 32].

Bacterial nitrogen fixation is the main mechanism that supplies nitrogen to holopelagic *Sargassum* in the GASB. The environmental characteristics of the open ocean promote rich diazotrophic biofilm, consequently reducing the potential nitrogen limitation factor through bacterial/seaweed interactions. Further investigations should focus on other nutrients, such as phosphorus, which may also play an important role in *Sargassum* proliferation [53, 54].

Our study identifies the crucial role of heterotrophic diazotrophs in the nitrogen cycle in the GASB. Our results suggest that this community contributes reactive nitrogen at a significant and basin-wide scale due to the area's vastness. Further research is necessary to investigate the complex interactions among diazotroph community, *Sargassum*, and the surrounding environment in order to fully understand nitrogen dynamics in this unique marine ecosystem.

Acknowledgements

We thank the Flotte Océanographique Française (FOF), Captain Pierre Samuel and the crew of ANTEA, Captain Jean Dumarais, and the crew of YERSIN. The authors acknowledge the support of Monaco-Explorations, the French Ministère de la Transition Écologique et Solidaire, IRD, CNES, and US191 IMAGO, which provided the logistics and implementation of the scientific equipment aboard the N/O ANTEA. The project has been supported by the OMICS, Platforms of the MIO.

Author contributions

Valérie Michotey, Sandrine Ruitton and Frédéric Ménard (designed the experiments). Valérie Michotey, Sandrine Ruitton, Sophie Guasco, Delphine Thibault, Thomas Changeux, Solène Connan, Valérie Stiger-Pouvreau and Thierry Thibaut (took part in the sampling). Sophie Guasco, Matéo Léger-Pigout, François Le Loc'h, and Jean-Marie Munaron (performed the experiments). Matéo Léger-Pigout and Elisabeth Navarro (analysed the raw sequencing data). Matéo Léger-Pigout, Valérie Michotey and Elisabeth Navarro (analysed the results). Matéo Léger-Pigout, Valérie Michotey and Elisabeth Navarro (wrote the paper with input to authors)

Supplementary material

Supplementary material is available at The ISME Journal online.

Conflicts of interest

The authors declare no competing interests.

Funding

The project leading to this publication has received funding from European FEDER Fund under project 1166-39417, ANR-19-SARG-0004 (ORIGINS), and ANR-19-SARG-0008 (SAVE-C).

Data availability

The nucleotide sequence data generated and analysed during the current study are available from the NCBI BioProject database

under the BioProject ID #PRJNA1017983. The data sets generated or analysed during this study are available in Supplementary Data 1–3.

References

- Ryther JH. The Sargasso Sea. Sci Am 1956;194:98–104. https:// doi.org/10.1038/scientificamerican0156-98. https://www.jstor. org/stable/24943833.
- Lapointe BE, West LE, Sutton TT et al. Ryther revisited: nutrient excretions by fishes enhance productivity of pelagic Sargassum in the western North Atlantic Ocean. J Exp Mar Biol Ecol 2014;458: 46–56. https://doi.org/10.1016/j.jembe.2014.05.002.
- Lapointe BE. A comparison of nutrient-limited productivity in Sargassum natans from neritic vs. oceanic waters of the western North Atlantic Ocean. Limnol Oceanogr 1995;40:625–33. https:// doi.org/10.4319/lo.1995.40.3.0625.
- Carpenter EJ. Nitrogen fixation by a blue-green epiphyte on pelagic Sargassum. Science 1972;178:1207–9. https://doi. org/10.1126/science.178.4066.1207.
- Phlips EJ, Willis M, Verchick A. Aspects of nitrogen fixation in Sargassum communities off the coast of Florida. J Exp Mar Biol Ecol 1986;102:99–119. https://doi.org/10.1016/0022-0981(86)90170-X.
- Johnson C, Dubbs LL, Piehler M. Reframing the contribution of pelagic Sargassum epiphytic N₂ fixation. PLoS One 2023; 18:e0289485. https://doi.org/10.1371/journal.pone.0289485.
- Baker P, Minzlaff U, Schoenle A et al. Potential contribution of surface-dwelling Sargassum algae to deep-sea ecosystems in the southern North Atlantic. Deep Sea Res Part II Top Stud Oceanogr 2018;148:21–34. https://doi.org/10.1016/j.dsr2.2017.10.002.
- Lapointe BE, Brewton RA, Herren LW et al. Nutrient content and stoichiometry of pelagic Sargassum reflects increasing nitrogen availability in the Atlantic Basin. Nat Commun 2021;12:3060. https://doi.org/10.1038/s41467-021-23135-7.
- Milledge JJ, Harvey PJ. Golden tides: problem or golden opportunity? The valorisation of Sargassum from beach inundations. J Mar Sci Eng 2016;4:60. https://doi.org/10.3390/jmse4030060.
- Thompson TM, Young BR, Baroutian S. Pelagic Sargassum for energy and fertiliser production in the Caribbean: a case study on Barbados. Renew Sust Energ Rev 2020;118:109564. https://doi. org/10.1016/j.rser.2019.109564.
- Salter MA, Rodríguez-Martínez RE, Álvarez-Filip L et al. Pelagic Sargassum as an emerging vector of high rate carbonate sediment import to tropical Atlantic coastlines. Glob Planet Change 2020;195:103332. https://doi.org/10.1016/j. gloplacha.2020.103332.
- Davis D, Simister R, Campbell S et al. Biomass composition of the golden tide pelagic seaweeds Sargassum fluitans and S. Natans (morphotypes I and VIII) to inform valorisation pathways. Sci Total Environ 2021;762:143134. https://doi.org/10.1016/ j.scitotenv.2020.143134.
- Rodríguez-Martínez RE, Medina-Valmaseda AE, Blanchon P et al. Faunal mortality associated with massive beaching and decomposition of pelagic Sargassum. Mar Pollut Bull 2019;146:201–5. https://doi.org/10.1016/j.marpolbul.2019.06.015.
- Maurer AS, Stapleton SP, Layman CA et al. The Atlantic Sargassum invasion impedes beach access for nesting sea turtles. Clim Change Ecol 2021;2:100034. https://doi.org/10.1016/j. ecochg.2021.100034.
- Maurer AS, Gross K, Stapleton SP. Beached Sargassum alters sand thermal environments: implications for incubating sea turtle eggs. J Exp Mar Biol Ecol 2022;546:151650. https://doi.org/10.1016/ j.jembe.2021.151650.

- Turner RE, Rabalais NN. Coastal eutrophication near the Mississippi river delta. Nature 1994;368:619–21. https://doi. org/10.1038/368619a0.
- 17. Gower J, Young E, King S. Satellite images suggest a new Sargassum source region in 2011. Remote Sens Lett 2013;4:764–73. https://doi.org/10.1080/2150704X.2013.796433.
- Abstracts from the 2013 Proceedings of the 65th Gulf and Caribbean Fisheries Institute (GCFI). D Johnson 2012;65:102–
 https://proceedings.gcfi.org/proceedings/the-sargassuminvasion-of-the-eastern-caribbean-and-dynamics-of-theequatorial-north-atlantic/.
- Wang M, Hu C, Barnes BB et al. The great Atlantic Sargassum belt. Science 2019;365:83–7. https://doi.org/10.1126/science.aaw7912.
- Resiere D, Mehdaoui H, Florentin J et al. Sargassum seaweed health menace in the Caribbean: clinical characteristics of a population exposed to hydrogen sulfide during the 2018 massive stranding. Clin Toxicol 2021;59:215-23. https://doi. org/10.1080/15563650.2020.1789162.
- Rodríguez-Martínez RE, Torres-Conde EG, Jordán-Dahlgren E. Pelagic Sargassum cleanup cost in Mexico. Ocean Coast Manag 2023;237:106542. https://doi.org/10.1016/j. ocecoaman.2023.106542.
- 22. Guidotti TL. Hydrogen sulphide. Occup Med 1996;**46**:367–71. https://doi.org/10.1093/occmed/46.5.367.
- Johns EM, Lumpkin R, Putman NF et al. The establishment of a pelagic Sargassum population in the tropical Atlantic: biological consequences of a basin-scale long distance dispersal event. Prog Oceanogr 2020;182:102269. https://doi.org/10.1016/j. pocean.2020.102269.
- Magaña-Gallegos E, García-Sánchez M, Graham C et al. Growth rates of pelagic Sargassum species in the Mexican Caribbean. Aquat Bot 2023;185:103614. https://doi.org/10.1016/ j.aquabot.2022.103614.
- Sissini MN, Barreto MBB D B, Széchy MTM et al. The floating Sargassum (Phaeophyceae) of the South Atlantic Ocean—likely scenarios. Phycologia 2017;56:321–8. https://doi.org/10.2216/ 16-92.1.
- Ward ND, Bianchi TS, Sawakuchi HO *et al*. The reactivity of plantderived organic matter and the potential importance of priming effects along the lower Amazon River. *J Geophys Res Biogeosci* 2016;**121**:1522–39. https://doi.org/10.1002/2016JG003342.
- Brooks MT, Coles VJ, Hood RR et al. Factors controlling the seasonal distribution of pelagic Sargassum. Mar Ecol Prog Ser 2018;599:1–18. https://doi.org/10.3354/meps12646.
- Berline L, Ody A, Jouanno J et al. Hindcasting the 2017 dispersal of Sargassum algae in the tropical North Atlantic. Mar Pollut Bull 2020;158:111431. https://doi.org/10.1016/j. marpolbul.2020.111431.
- Jouanno J, Moquet JS, Berline L et al. Evolution of the riverine nutrient export to the tropical Atlantic over the last 15 years: is there a link with Sargassum proliferation? Environ Res Lett 2021;16:034042. https://doi.org/10.1088/1748-9326/abe11a.
- Michotey V, Blanfuné A, Chevalier C et al. In situ observations and modelling revealed environmental factors favouring occurrence of vibrio in microbiome of the pelagic Sargassum responsible for strandings. Sci Total Environ 2020;**748**:141216. https://doi. org/10.1016/j.scitotenv.2020.141216.
- Torralba MG, Franks JS, Gomez A et al. Effect of Macondo Prospect 252 oil on microbiota associated with pelagic Sargassum in the northern Gulf of Mexico. Microb Ecol 2017;**73**:91–100. https://doi.org/10.1007/s00248-016-0857-y.
- 32. Theirlynck T, Mendonça IRW, Engelen AH et al. Diversity of the holopelagic Sargassum microbiome from the great Atlantic

Sargassum Belt to coastal stranding locations. Harmful Algae 2023;**122**:102369. https://doi.org/10.1016/j.hal.2022.102369.

- Schell J, Goodwin D, Siuda A. Recent Sargassum inundation events in the Caribbean: shipboard observations reveal dominance of a previously rare form. Oceanography 2015;28:8–10. https://doi.org/10.5670/oceanog.2015.70.
- 34. Ody A, Thibaut T, Berline L et al. From in situ to satellite observations of pelagic Sargassum distribution and aggregation in the tropical North Atlantic Ocean. PLoS One 2019;14:e0222584. https://doi.org/10.1371/journal.pone.0222584.
- Henry S, Bru D, Stres B et al. Quantitative detection of the nosZ gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, narG, nirK, and nosZ genes in soils. Appl Environ Microbiol 2006;**72**:5181–9. https://doi.org/10.1128/ AEM.00231-06.
- Rösch C, Mergel A, Bothe H. Biodiversity of denitrifying and dinitrogen-fixing bacteria in an acid Forest soil. Appl Environ Microbiol 2002;68:3818–29. https://doi.org/10.1128/ AEM.68.8.3818-3829.2002.
- Rotthauwe JH, Witzel KP, Liesack W. The ammonia monooxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Appl Environ Microbiol* 1997;**63**:4704. 4712, https://doi. org/10.1128/aem.63.12.4704-4712.1997.
- Zaghmouri I, Michotey VD, Guasco S et al. Trends in salinity and inorganic nitrogen compounds in the Berre lagoon (1994–2011) bacterial activities and nitrogen budgets. Mar Pollut Bull 2013;67: 55–65. https://doi.org/10.1016/j.marpolbul.2012.11.046.
- Zehr JP, Turner PJ. Nitrogen fixation: nitrogenase genes and gene expression. Methods Microbiol 2001;30:271–86. https://doi. org/10.1016/S0580-9517(01)30049-1.
- Poly F, Monrozier LJ, Bally R. Improvement in the RFLP procedure for studying the diversity of nifH genes in communities of nitrogen fixers in soil. Res Microbiol 2001;152:95–103. https://doi. org/10.1016/S0923-2508(00)01172-4.
- Gaby JC, Buckley DH. A comprehensive evaluation of PCR primers to amplify the *nifH* gene of nitrogenase. *PLoS One* 2012;**7**:e42149. https://doi.org/10.1371/journal. pone.0042149.
- Callahan BJ, McMurdie PJ, Rosen MJ et al. DADA2: high-resolution sample inference from Illumina amplicon data. Nat Methods 2016;13:581–3. https://doi.org/10.1038/nmeth.3869.
- Heller P, Tripp HJ, Turk-Kubo K et al. ARBitrator: a software pipeline for on-demand retrieval of auto-curated nifH sequences from GenBank. Bioinformatics 2014;30:2883–90. https://doi. org/10.1093/bioinformatics/btu417.
- 44. Hanson RB. Pelagic Sargassum community metabolism: carbon and nitrogen. J Exp Mar Biol Ecol 1977;29:107–18. https://doi. org/10.1016/0022-0981(77)90042-9.
- 45. Phlips E, Zeman C. Photosynthesis, growth and nitrogen fixation by epiphytic forms of filamentous cyanobacteria from pelagic Sargassum. Bull Mar Sci 1990;47:613–21. https:// www.ingentaconnect.com/content/umrsmas/bullmar/1990/00 000047/00000003/art00003.
- Rooker JR, Turner JP, Holt SA. Trophic ecology of Sargassumassociated fishes in the Gulf of Mexico determined from stable isotopes and fatty acids. Mar Ecol Prog Ser 2006;**313**:249–59. https://doi.org/10.3354/meps313249.
- Vázquez-Delfín E, Freile-Pelegrín Y, Salazar-Garibay A et al. Species composition and chemical characterization of Sargassum influx at six different locations along the Mexican Caribbean coast. Sci Total Environ 2021;795:148852. https://doi. org/10.1016/j.scitotenv.2021.148852.

- Hervé V, Lambourdière J, René-Trouillefou M et al. Sargassum differentially shapes the microbiota composition and diversity at coastal tide sites and inland storage sites on Caribbean Islands. Front Microbiol 2021;12:3221. https://doi.org/10.3389/ fmicb.2021.701155.
- Mohapatra BR. Phylogenetic and functional characterization of the microbiome of Sargassum seaweed waste. Appl Phycol 2023;4:87–98. https://doi.org/10.1080/26388081.2023. 2220384.
- Montoya JP. Nitrogen stable isotopes in marine environments. In: Douglas G Capone, Deborah A Bronk, Margaret R Mulholland, Edward J Carpenter, Nitrogen in the Marine Environment (Second Edition), Academic Press, London. https://doi.org/10.1016/ B978-0-12-372522-6.00029-3.
- Brylinsky M. Release of dissolved organic matter by some marine macrophytes. Mar Biol 1977;39:213–20. https://doi.org/10.1007/ BF00390995.
- Powers LC, Hertkorn N, McDonald N et al. Sargassum sp. act as a large regional source of marine dissolved organic carbon and polyphenols. Glob Biogeochem Cycles 2019;33:1423–39. https://doi. org/10.1029/2019GB006225.
- Pierre WH. The phosphorus cycle and soil fertility 1. Agron J 1948;40:1–14. https://doi.org/10.2134/agronj1948.00021962004 000010001x.
- 54. Ohtake M, Natori N, Sugai Y et al. Growth and nutrient uptake characteristics of *Sargassum macrocarpum* cultivated with phosphorus-replete wastewater. Aquat Bot 2020;**163**:103208. https://doi.org/10.1016/j.aquabot.2020.103208.